

P2.1 Position — Time

An object's position can be measured and graphed as a function of time. An object's speed can be calculated and graphed as a function of time.
molecules that make up living systems:

P2.1 Position — Time

An object's position can be measured and graphed as a function of time. An object's speed can be calculated and graphed as a function of time.
molecules that make up living systems:

P2.1 Position — Time

An object's position can be measured and graphed as a function of time. An object's speed can be calculated and graphed as a function of time.
molecules that make up living systems:

P2.1 Position — Time

An object's position can be measured and graphed as a function of time. An object's speed can be calculated and graphed as a function of time.
molecules that make up living systems:

P2.1 Position — Time

An object's position can be measured and graphed as a function of time. An object's speed can be calculated and graphed as a function of time.
molecules that make up living systems:

P2.1 Position — Time

An object's position can be measured and graphed as a function of time. An object's speed can be calculated and graphed as a function of time.
molecules that make up living systems:

E

P2.1A Calculate the average speed of an object using the change of position and elapsed time.

E

P2.1A Calculate the average speed of an object using the change of position and elapsed time.

E

P2.1A Calculate the average speed of an object using the change of position and elapsed time.

E

P2.1A Calculate the average speed of an object using the change of position and elapsed time.

E

P2.1A Calculate the average speed of an object using the change of position and elapsed time.

E

P2.1A Calculate the average speed of an object using the change of position and elapsed time.

P2.1 Position — Time

An object's position can be measured and graphed as a function of time. An object's speed can be calculated and graphed as a function of time.

P2.1 Position — Time

An object's position can be measured and graphed as a function of time. An object's speed can be calculated and graphed as a function of time.

P2.1 Position — Time

An object's position can be measured and graphed as a function of time. An object's speed can be calculated and graphed as a function of time.

P2.1 Position — Time

An object's position can be measured and graphed as a function of time. An object's speed can be calculated and graphed as a function of time.

P2.1 Position — Time

An object's position can be measured and graphed as a function of time. An object's speed can be calculated and graphed as a function of time.

P2.1 Position — Time

An object's position can be measured and graphed as a function of time. An object's speed can be calculated and graphed as a function of time.



P2.1B Represent the velocities for linear and circular motion using motion diagrams (arrows on strobe pictures).



P2.1B Represent the velocities for linear and circular motion using motion diagrams (arrows on strobe pictures).



P2.1B Represent the velocities for linear and circular motion using motion diagrams (arrows on strobe pictures).



P2.1B Represent the velocities for linear and circular motion using motion diagrams (arrows on strobe pictures).



P2.1B Represent the velocities for linear and circular motion using motion diagrams (arrows on strobe pictures).



P2.1B Represent the velocities for linear and circular motion using motion diagrams (arrows on strobe pictures).

P2.1 Position — Time

An object's position can be measured and graphed as a function of time. An object's speed can be calculated and graphed as a function of time.

P2.1 Position — Time

An object's position can be measured and graphed as a function of time. An object's speed can be calculated and graphed as a function of time.

P2.1 Position — Time

An object's position can be measured and graphed as a function of time. An object's speed can be calculated and graphed as a function of time.

P2.1 Position — Time

An object's position can be measured and graphed as a function of time. An object's speed can be calculated and graphed as a function of time.

P2.1 Position — Time

An object's position can be measured and graphed as a function of time. An object's speed can be calculated and graphed as a function of time.

P2.1 Position — Time

An object's position can be measured and graphed as a function of time. An object's speed can be calculated and graphed as a function of time.



P2.IC Create line graphs using measured values of position and elapsed time.



P2.IC Create line graphs using measured values of position and elapsed time.



P2.IC Create line graphs using measured values of position and elapsed time.



P2.IC Create line graphs using measured values of position and elapsed time.



P2.IC Create line graphs using measured values of position and elapsed time.



P2.IC Create line graphs using measured values of position and elapsed time.

P2.1 Position — Time

An object's position can be measured and graphed as a function of time. An object's speed can be calculated and graphed as a function of time.

P2.1 Position — Time

An object's position can be measured and graphed as a function of time. An object's speed can be calculated and graphed as a function of time.

P2.1 Position — Time

An object's position can be measured and graphed as a function of time. An object's speed can be calculated and graphed as a function of time.

P2.1 Position — Time

An object's position can be measured and graphed as a function of time. An object's speed can be calculated and graphed as a function of time.

P2.1 Position — Time

An object's position can be measured and graphed as a function of time. An object's speed can be calculated and graphed as a function of time.

P2.1 Position — Time

An object's position can be measured and graphed as a function of time. An object's speed can be calculated and graphed as a function of time.

E

P2.ID Describe and analyze the motion that a position-time graph represents, given the graph.

E

P2.ID Describe and analyze the motion that a position-time graph represents, given the graph.

E

P2.ID Describe and analyze the motion that a position-time graph represents, given the graph.

E

P2.ID Describe and analyze the motion that a position-time graph represents, given the graph.

E

P2.ID Describe and analyze the motion that a position-time graph represents, given the graph.

E

P2.ID Describe and analyze the motion that a position-time graph represents, given the graph.

P2.1 Position — Time

An object's position can be measured and graphed as a function of time. An object's speed can be calculated and graphed as a function of time.

P2.1 Position — Time

An object's position can be measured and graphed as a function of time. An object's speed can be calculated and graphed as a function of time.

P2.1 Position — Time

An object's position can be measured and graphed as a function of time. An object's speed can be calculated and graphed as a function of time.

P2.1 Position — Time

An object's position can be measured and graphed as a function of time. An object's speed can be calculated and graphed as a function of time.

P2.1 Position — Time

An object's position can be measured and graphed as a function of time. An object's speed can be calculated and graphed as a function of time.

P2.1 Position — Time

An object's position can be measured and graphed as a function of time. An object's speed can be calculated and graphed as a function of time.



P2.1E Describe and classify various motions in a plane as one dimensional, two dimensional, circular, or periodic.



P2.1E Describe and classify various motions in a plane as one dimensional, two dimensional, circular, or periodic.



P2.1E Describe and classify various motions in a plane as one dimensional, two dimensional, circular, or periodic.



P2.1E Describe and classify various motions in a plane as one dimensional, two dimensional, circular, or periodic.



PP2.1E Describe and classify various motions in a plane as one dimensional, two dimensional, circular, or periodic.



P2.1E Describe and classify various motions in a plane as one dimensional, two dimensional, circular, or periodic.

P2.1 Position — Time

An object's position can be measured and graphed as a function of time. An object's speed can be calculated and graphed as a function of time.

P2.1 Position — Time

An object's position can be measured and graphed as a function of time. An object's speed can be calculated and graphed as a function of time.

P2.1 Position — Time

An object's position can be measured and graphed as a function of time. An object's speed can be calculated and graphed as a function of time.

P2.1 Position — Time

An object's position can be measured and graphed as a function of time. An object's speed can be calculated and graphed as a function of time.

P2.1 Position — Time

An object's position can be measured and graphed as a function of time. An object's speed can be calculated and graphed as a function of time.

P2.1 Position — Time

An object's position can be measured and graphed as a function of time. An object's speed can be calculated and graphed as a function of time.

E

P2.IF Distinguish between rotation and revolution and describe and contrast the two speeds of an object like the Earth.

E

P2.IF Distinguish between rotation and revolution and describe and contrast the two speeds of an object like the Earth.

E

P2.IF Distinguish between rotation and revolution and describe and contrast the two speeds of an object like the Earth.

E

P2.IF Distinguish between rotation and revolution and describe and contrast the two speeds of an object like the Earth.

E

P2.IF Distinguish between rotation and revolution and describe and contrast the two speeds of an object like the Earth.

E

P2.IF Distinguish between rotation and revolution and describe and contrast the two speeds of an object like the Earth.

P2.1 Position — Time

An object's position can be measured and graphed as a function of time. An object's speed can be calculated and graphed as a function of time.

P2.1 Position — Time

An object's position can be measured and graphed as a function of time. An object's speed can be calculated and graphed as a function of time.

P2.1 Position — Time

An object's position can be measured and graphed as a function of time. An object's speed can be calculated and graphed as a function of time.

P2.1 Position — Time

An object's position can be measured and graphed as a function of time. An object's speed can be calculated and graphed as a function of time.

P2.1 Position — Time

An object's position can be measured and graphed as a function of time. An object's speed can be calculated and graphed as a function of time.

P2.1 Position — Time

An object's position can be measured and graphed as a function of time. An object's speed can be calculated and graphed as a function of time.



P2.1g Solve problems involving average speed and constant acceleration in one dimension.



P2.1g Solve problems involving average speed and constant acceleration in one dimension.



P2.1g Solve problems involving average speed and constant acceleration in one dimension.



P2.1g Solve problems involving average speed and constant acceleration in one dimension.



P2.1g Solve problems involving average speed and constant acceleration in one dimension.



P2.1g Solve problems involving average speed and constant acceleration in one dimension.

P2.1 Position — Time

An object's position can be measured and graphed as a function of time. An object's speed can be calculated and graphed as a function of time.

P2.1 Position — Time

An object's position can be measured and graphed as a function of time. An object's speed can be calculated and graphed as a function of time.

P2.1 Position — Time

An object's position can be measured and graphed as a function of time. An object's speed can be calculated and graphed as a function of time.

P2.1 Position — Time

An object's position can be measured and graphed as a function of time. An object's speed can be calculated and graphed as a function of time.

P2.1 Position — Time

An object's position can be measured and graphed as a function of time. An object's speed can be calculated and graphed as a function of time.

P2.1 Position — Time

An object's position can be measured and graphed as a function of time. An object's speed can be calculated and graphed as a function of time.



P2.1h Identify the changes in speed and direction in everyday examples of circular (rotation and revolution), periodic, and projectile motions.



P2.1h Identify the changes in speed and direction in everyday examples of circular (rotation and revolution), periodic, and projectile motions.



P2.1h Identify the changes in speed and direction in everyday examples of circular (rotation and revolution), periodic, and projectile motions.



P2.1h Identify the changes in speed and direction in everyday examples of circular (rotation and revolution), periodic, and projectile motions.



P2.1h Identify the changes in speed and direction in everyday examples of circular (rotation and revolution), periodic, and projectile motions.



P2.1h Identify the changes in speed and direction in everyday examples of circular (rotation and revolution), periodic, and projectile motions.

P2.2 Velocity — Time

The motion of an object can be described by its position and velocity as functions of time and by its average speed and average acceleration during intervals of time.

P2.2 Velocity — Time

The motion of an object can be described by its position and velocity as functions of time and by its average speed and average acceleration during intervals of time.

P2.2 Velocity — Time

The motion of an object can be described by its position and velocity as functions of time and by its average speed and average acceleration during intervals of time.

P2.2 Velocity — Time

The motion of an object can be described by its position and velocity as functions of time and by its average speed and average acceleration during intervals of time.

P2.2 Velocity — Time

The motion of an object can be described by its position and velocity as functions of time and by its average speed and average acceleration during intervals of time.

P2.2 Velocity — Time

The motion of an object can be described by its position and velocity as functions of time and by its average speed and average acceleration during intervals of time.



P2.2A Distinguish between the variables of distance, displacement, speed, velocity, and acceleration.



P2.2A Distinguish between the variables of distance, displacement, speed, velocity, and acceleration.



P2.2A Distinguish between the variables of distance, displacement, speed, velocity, and acceleration.



P2.2A Distinguish between the variables of distance, displacement, speed, velocity, and acceleration.



P2.2A Distinguish between the variables of distance, displacement, speed, velocity, and acceleration.



P2.2A Distinguish between the variables of distance, displacement, speed, velocity, and acceleration.

P2.2 Velocity — Time

The motion of an object can be described by its position and velocity as functions of time and by its average speed and average acceleration during intervals of time.

P2.2 Velocity — Time

The motion of an object can be described by its position and velocity as functions of time and by its average speed and average acceleration during intervals of time.

P2.2 Velocity — Time

The motion of an object can be described by its position and velocity as functions of time and by its average speed and average acceleration during intervals of time.

P2.2 Velocity — Time

The motion of an object can be described by its position and velocity as functions of time and by its average speed and average acceleration during intervals of time.

P2.2 Velocity — Time

The motion of an object can be described by its position and velocity as functions of time and by its average speed and average acceleration during intervals of time.

P2.2 Velocity — Time

The motion of an object can be described by its position and velocity as functions of time and by its average speed and average acceleration during intervals of time.



P2.2B Use the change of speed and elapsed time to calculate the average acceleration for linear motion.



P2.2B Use the change of speed and elapsed time to calculate the average acceleration for linear motion.



P2.2B Use the change of speed and elapsed time to calculate the average acceleration for linear motion.



P2.2B Use the change of speed and elapsed time to calculate the average acceleration for linear motion.



P2.2B Use the change of speed and elapsed time to calculate the average acceleration for linear motion.



P2.2B Use the change of speed and elapsed time to calculate the average acceleration for linear motion.

P2.2 Velocity — Time

The motion of an object can be described by its position and velocity as functions of time and by its average speed and average acceleration during intervals of time.

P2.2 Velocity — Time

The motion of an object can be described by its position and velocity as functions of time and by its average speed and average acceleration during intervals of time.

P2.2 Velocity — Time

The motion of an object can be described by its position and velocity as functions of time and by its average speed and average acceleration during intervals of time.

P2.2 Velocity — Time

The motion of an object can be described by its position and velocity as functions of time and by its average speed and average acceleration during intervals of time.

P2.2 Velocity — Time

The motion of an object can be described by its position and velocity as functions of time and by its average speed and average acceleration during intervals of time.

P2.2 Velocity — Time

The motion of an object can be described by its position and velocity as functions of time and by its average speed and average acceleration during intervals of time.

E

P2.2C Describe and analyze the motion that a velocity-time graph represents, given the graph.

E

P2.2C Describe and analyze the motion that a velocity-time graph represents, given the graph.

E

P2.2C Describe and analyze the motion that a velocity-time graph represents, given the graph.

E

P2.2C Describe and analyze the motion that a velocity-time graph represents, given the graph.

E

P2.2C Describe and analyze the motion that a velocity-time graph represents, given the graph.

E

P2.2C Describe and analyze the motion that a velocity-time graph represents, given the graph.

P2.2 Velocity — Time

The motion of an object can be described by its position and velocity as functions of time and by its average speed and average acceleration during intervals of time.

P2.2 Velocity — Time

The motion of an object can be described by its position and velocity as functions of time and by its average speed and average acceleration during intervals of time.

P2.2 Velocity — Time

The motion of an object can be described by its position and velocity as functions of time and by its average speed and average acceleration during intervals of time.

P2.2 Velocity — Time

The motion of an object can be described by its position and velocity as functions of time and by its average speed and average acceleration during intervals of time.

P2.2 Velocity — Time

The motion of an object can be described by its position and velocity as functions of time and by its average speed and average acceleration during intervals of time.

P2.2 Velocity — Time

The motion of an object can be described by its position and velocity as functions of time and by its average speed and average acceleration during intervals of time.



P2.2D State that uniform circular motion involves acceleration without a change in speed.



P2.2D State that uniform circular motion involves acceleration without a change in speed.



P2.2D State that uniform circular motion involves acceleration without a change in speed.



P2.2D State that uniform circular motion involves acceleration without a change in speed.



P2.2D State that uniform circular motion involves acceleration without a change in speed.



P2.2D State that uniform circular motion involves acceleration without a change in speed.

P2.2 Velocity — Time

The motion of an object can be described by its position and velocity as functions of time and by its average speed and average acceleration during intervals of time.

P2.2 Velocity — Time

The motion of an object can be described by its position and velocity as functions of time and by its average speed and average acceleration during intervals of time.

P2.2 Velocity — Time

The motion of an object can be described by its position and velocity as functions of time and by its average speed and average acceleration during intervals of time.

P2.2 Velocity — Time

The motion of an object can be described by its position and velocity as functions of time and by its average speed and average acceleration during intervals of time.

P2.2 Velocity — Time

The motion of an object can be described by its position and velocity as functions of time and by its average speed and average acceleration during intervals of time.

P2.2 Velocity — Time

The motion of an object can be described by its position and velocity as functions of time and by its average speed and average acceleration during intervals of time.



P2.2e Use the area under a velocity-time graph to calculate the distance traveled and the slope to calculate the acceleration.



P2.2e Use the area under a velocity-time graph to calculate the distance traveled and the slope to calculate the acceleration.



P2.2e Use the area under a velocity-time graph to calculate the distance traveled and the slope to calculate the acceleration.



P2.2e Use the area under a velocity-time graph to calculate the distance traveled and the slope to calculate the acceleration.



P2.2e Use the area under a velocity-time graph to calculate the distance traveled and the slope to calculate the acceleration.



P2.2e Use the area under a velocity-time graph to calculate the distance traveled and the slope to calculate the acceleration.

P2.2 Velocity — Time

The motion of an object can be described by its position and velocity as functions of time and by its average speed and average acceleration during intervals of time.

P2.2 Velocity — Time

The motion of an object can be described by its position and velocity as functions of time and by its average speed and average acceleration during intervals of time.

P2.2 Velocity — Time

The motion of an object can be described by its position and velocity as functions of time and by its average speed and average acceleration during intervals of time.

P2.2 Velocity — Time

The motion of an object can be described by its position and velocity as functions of time and by its average speed and average acceleration during intervals of time.

P2.2 Velocity — Time

The motion of an object can be described by its position and velocity as functions of time and by its average speed and average acceleration during intervals of time.

P2.2 Velocity — Time

The motion of an object can be described by its position and velocity as functions of time and by its average speed and average acceleration during intervals of time.



P2.2f Describe the relationship between changes in position, velocity, and acceleration during periodic motion.



P2.2f Describe the relationship between changes in position, velocity, and acceleration during periodic motion.



P2.2f Describe the relationship between changes in position, velocity, and acceleration during periodic motion.



P2.2f Describe the relationship between changes in position, velocity, and acceleration during periodic motion.



P2.2f Describe the relationship between changes in position, velocity, and acceleration during periodic motion.



P2.2f Describe the relationship between changes in position, velocity, and acceleration during periodic motion.

P2.2 Velocity — Time

The motion of an object can be described by its position and velocity as functions of time and by its average speed and average acceleration during intervals of time.

P2.2 Velocity — Time

The motion of an object can be described by its position and velocity as functions of time and by its average speed and average acceleration during intervals of time.

P2.2 Velocity — Time

The motion of an object can be described by its position and velocity as functions of time and by its average speed and average acceleration during intervals of time.

P2.2 Velocity — Time

The motion of an object can be described by its position and velocity as functions of time and by its average speed and average acceleration during intervals of time.

P2.2 Velocity — Time

The motion of an object can be described by its position and velocity as functions of time and by its average speed and average acceleration during intervals of time.

P2.2 Velocity — Time

The motion of an object can be described by its position and velocity as functions of time and by its average speed and average acceleration during intervals of time.



P2.2g Apply the independence of the vertical and horizontal initial velocities to solve projectile motion problems.



P2.2g Apply the independence of the vertical and horizontal initial velocities to solve projectile motion problems.



P2.2g Apply the independence of the vertical and horizontal initial velocities to solve projectile motion problems.



P2.2g Apply the independence of the vertical and horizontal initial velocities to solve projectile motion problems.



P2.2g Apply the independence of the vertical and horizontal initial velocities to solve projectile motion problems.



P2.2g Apply the independence of the vertical and horizontal initial velocities to solve projectile motion problems.

P2.3x Frames of Reference

All motion is relative to whatever frame of reference is chosen, for there is no motionless frame from which to judge all motion.

P2.3x Frames of Reference

All motion is relative to whatever frame of reference is chosen, for there is no motionless frame from which to judge all motion.

P2.3x Frames of Reference

All motion is relative to whatever frame of reference is chosen, for there is no motionless frame from which to judge all motion.

P2.3x Frames of Reference

All motion is relative to whatever frame of reference is chosen, for there is no motionless frame from which to judge all motion.

P2.3x Frames of Reference

All motion is relative to whatever frame of reference is chosen, for there is no motionless frame from which to judge all motion.

P2.3x Frames of Reference

All motion is relative to whatever frame of reference is chosen, for there is no motionless frame from which to judge all motion.



P2.3a Describe and compare the motion of an object using different reference frames.



P2.3a Describe and compare the motion of an object using different reference frames.



P2.3a Describe and compare the motion of an object using different reference frames.



P2.3a Describe and compare the motion of an object using different reference frames.



P2.3a Describe and compare the motion of an object using different reference frames.



P2.3a Describe and compare the motion of an object using different reference frames.

P3.1 Basic Forces in Nature

Objects can interact with each other by “direct contact” (e.g., pushes or pulls, friction) or at a distance (e.g., gravity, electromagnetism, nuclear).

P3.1 Basic Forces in Nature

Objects can interact with each other by “direct contact” (e.g., pushes or pulls, friction) or at a distance (e.g., gravity, electromagnetism, nuclear).

P3.1 Basic Forces in Nature

Objects can interact with each other by “direct contact” (e.g., pushes or pulls, friction) or at a distance (e.g., gravity, electromagnetism, nuclear).

P3.1 Basic Forces in Nature

Objects can interact with each other by “direct contact” (e.g., pushes or pulls, friction) or at a distance (e.g., gravity, electromagnetism, nuclear).

P3.1 Basic Forces in Nature

Objects can interact with each other by “direct contact” (e.g., pushes or pulls, friction) or at a distance (e.g., gravity, electromagnetism, nuclear).

P3.1 Basic Forces in Nature

Objects can interact with each other by “direct contact” (e.g., pushes or pulls, friction) or at a distance (e.g., gravity, electromagnetism, nuclear).



P3.1A Identify the force(s) acting between objects in “direct contact” or at a distance.



P3.1A Identify the force(s) acting between objects in “direct contact” or at a distance.



P3.1A Identify the force(s) acting between objects in “direct contact” or at a distance.



P3.1A Identify the force(s) acting between objects in “direct contact” or at a distance.



P3.1A Identify the force(s) acting between objects in “direct contact” or at a distance.



P3.1A Identify the force(s) acting between objects in “direct contact” or at a distance.

P3.1x Forces

There are four basic forces (gravitational, electromagnetic, strong, and weak nuclear) that differ greatly in magnitude and range. Between any two charged particles, electric force is vastly greater than the gravitational force. Most observable forces (e.g., those exerted by a coiled spring or friction) may be traced to electric forces acting between atoms and molecules.

P3.1x Forces

There are four basic forces (gravitational, electromagnetic, strong, and weak nuclear) that differ greatly in magnitude and range. Between any two charged particles, electric force is vastly greater than the gravitational force. Most observable forces (e.g., those exerted by a coiled spring or friction) may be traced to electric forces acting between atoms and molecules.

P3.1x Forces

There are four basic forces (gravitational, electromagnetic, strong, and weak nuclear) that differ greatly in magnitude and range. Between any two charged particles, electric force is vastly greater than the gravitational force. Most observable forces (e.g., those exerted by a coiled spring or friction) may be traced to electric forces acting between atoms and molecules.

P3.1x Forces

There are four basic forces (gravitational, electromagnetic, strong, and weak nuclear) that differ greatly in magnitude and range. Between any two charged particles, electric force is vastly greater than the gravitational force. Most observable forces (e.g., those exerted by a coiled spring or friction) may be traced to electric forces acting between atoms and molecules.

P3.1x Forces

There are four basic forces (gravitational, electromagnetic, strong, and weak nuclear) that differ greatly in magnitude and range. Between any two charged particles, electric force is vastly greater than the gravitational force. Most observable forces (e.g., those exerted by a coiled spring or friction) may be traced to electric forces acting between atoms and molecules.

P3.1x Forces

There are four basic forces (gravitational, electromagnetic, strong, and weak nuclear) that differ greatly in magnitude and range. Between any two charged particles, electric force is vastly greater than the gravitational force. Most observable forces (e.g., those exerted by a coiled spring or friction) may be traced to electric forces acting between atoms and molecules.



P3.1b Explain why scientists can ignore the gravitational force when measuring the net force between two electrons.



P3.1b Explain why scientists can ignore the gravitational force when measuring the net force between two electrons.



P3.1b Explain why scientists can ignore the gravitational force when measuring the net force between two electrons.



P3.1b Explain why scientists can ignore the gravitational force when measuring the net force between two electrons.



P3.1b Explain why scientists can ignore the gravitational force when measuring the net force between two electrons.



P3.1b Explain why scientists can ignore the gravitational force when measuring the net force between two electrons.

P3.1x Forces

There are four basic forces (gravitational, electromagnetic, strong, and weak nuclear) that differ greatly in magnitude and range. Between any two charged particles, electric force is vastly greater than the gravitational force. Most observable forces (e.g., those exerted by a coiled spring or friction) may be traced to electric forces acting between atoms and molecules.

P3.1x Forces

There are four basic forces (gravitational, electromagnetic, strong, and weak nuclear) that differ greatly in magnitude and range. Between any two charged particles, electric force is vastly greater than the gravitational force. Most observable forces (e.g., those exerted by a coiled spring or friction) may be traced to electric forces acting between atoms and molecules.

P3.1x Forces

There are four basic forces (gravitational, electromagnetic, strong, and weak nuclear) that differ greatly in magnitude and range. Between any two charged particles, electric force is vastly greater than the gravitational force. Most observable forces (e.g., those exerted by a coiled spring or friction) may be traced to electric forces acting between atoms and molecules.

P3.1x Forces

There are four basic forces (gravitational, electromagnetic, strong, and weak nuclear) that differ greatly in magnitude and range. Between any two charged particles, electric force is vastly greater than the gravitational force. Most observable forces (e.g., those exerted by a coiled spring or friction) may be traced to electric forces acting between atoms and molecules.

P3.1x Forces

There are four basic forces (gravitational, electromagnetic, strong, and weak nuclear) that differ greatly in magnitude and range. Between any two charged particles, electric force is vastly greater than the gravitational force. Most observable forces (e.g., those exerted by a coiled spring or friction) may be traced to electric forces acting between atoms and molecules.

P3.1x Forces

There are four basic forces (gravitational, electromagnetic, strong, and weak nuclear) that differ greatly in magnitude and range. Between any two charged particles, electric force is vastly greater than the gravitational force. Most observable forces (e.g., those exerted by a coiled spring or friction) may be traced to electric forces acting between atoms and molecules.



P3.1c Provide examples that illustrate the importance of the electric force in everyday life.



P3.1c Provide examples that illustrate the importance of the electric force in everyday life.



P3.1c Provide examples that illustrate the importance of the electric force in everyday life.



P3.1c Provide examples that illustrate the importance of the electric force in everyday life.



P3.1c Provide examples that illustrate the importance of the electric force in everyday life.



P3.1c Provide examples that illustrate the importance of the electric force in everyday life.

P3.1x Forces

There are four basic forces (gravitational, electromagnetic, strong, and weak nuclear) that differ greatly in magnitude and range. Between any two charged particles, electric force is vastly greater than the gravitational force. Most observable forces (e.g., those exerted by a coiled spring or friction) may be traced to electric forces acting between atoms and molecules.

P3.1x Forces

There are four basic forces (gravitational, electromagnetic, strong, and weak nuclear) that differ greatly in magnitude and range. Between any two charged particles, electric force is vastly greater than the gravitational force. Most observable forces (e.g., those exerted by a coiled spring or friction) may be traced to electric forces acting between atoms and molecules.

P3.1x Forces

There are four basic forces (gravitational, electromagnetic, strong, and weak nuclear) that differ greatly in magnitude and range. Between any two charged particles, electric force is vastly greater than the gravitational force. Most observable forces (e.g., those exerted by a coiled spring or friction) may be traced to electric forces acting between atoms and molecules.

P3.1x Forces

There are four basic forces (gravitational, electromagnetic, strong, and weak nuclear) that differ greatly in magnitude and range. Between any two charged particles, electric force is vastly greater than the gravitational force. Most observable forces (e.g., those exerted by a coiled spring or friction) may be traced to electric forces acting between atoms and molecules.

P3.1x Forces

There are four basic forces (gravitational, electromagnetic, strong, and weak nuclear) that differ greatly in magnitude and range. Between any two charged particles, electric force is vastly greater than the gravitational force. Most observable forces (e.g., those exerted by a coiled spring or friction) may be traced to electric forces acting between atoms and molecules.

P3.1x Forces

There are four basic forces (gravitational, electromagnetic, strong, and weak nuclear) that differ greatly in magnitude and range. Between any two charged particles, electric force is vastly greater than the gravitational force. Most observable forces (e.g., those exerted by a coiled spring or friction) may be traced to electric forces acting between atoms and molecules.



P3.Id Identify the basic forces in everyday interactions.



P3.Id Identify the basic forces in everyday interactions.



P3.Id Identify the basic forces in everyday interactions.



P3.Id Identify the basic forces in everyday interactions.



P3.Id Identify the basic forces in everyday interactions.



P3.Id Identify the basic forces in everyday interactions.

P3.2 Net Forces

Forces have magnitude and direction. The net force on an object is the sum of all the forces acting on the object. Objects change their speed and/or direction only when a net force is applied. If the net force on an object is zero, there is no change in motion (Newton's First Law).

P3.2 Net Forces

Forces have magnitude and direction. The net force on an object is the sum of all the forces acting on the object. Objects change their speed and/or direction only when a net force is applied. If the net force on an object is zero, there is no change in motion (Newton's First Law).

P3.2 Net Forces

Forces have magnitude and direction. The net force on an object is the sum of all the forces acting on the object. Objects change their speed and/or direction only when a net force is applied. If the net force on an object is zero, there is no change in motion (Newton's First Law).

P3.2 Net Forces

Forces have magnitude and direction. The net force on an object is the sum of all the forces acting on the object. Objects change their speed and/or direction only when a net force is applied. If the net force on an object is zero, there is no change in motion (Newton's First Law).

P3.2 Net Forces

Forces have magnitude and direction. The net force on an object is the sum of all the forces acting on the object. Objects change their speed and/or direction only when a net force is applied. If the net force on an object is zero, there is no change in motion (Newton's First Law).

P3.2 Net Forces

Forces have magnitude and direction. The net force on an object is the sum of all the forces acting on the object. Objects change their speed and/or direction only when a net force is applied. If the net force on an object is zero, there is no change in motion (Newton's First Law).



P3.2A Identify the magnitude and direction of everyday forces (e.g., wind, tension in ropes, pushes and pulls, weight).



P3.2A Identify the magnitude and direction of everyday forces (e.g., wind, tension in ropes, pushes and pulls, weight).



P3.2A Identify the magnitude and direction of everyday forces (e.g., wind, tension in ropes, pushes and pulls, weight).



P3.2A Identify the magnitude and direction of everyday forces (e.g., wind, tension in ropes, pushes and pulls, weight).



P3.2A Identify the magnitude and direction of everyday forces (e.g., wind, tension in ropes, pushes and pulls, weight).



P3.2A Identify the magnitude and direction of everyday forces (e.g., wind, tension in ropes, pushes and pulls, weight).

P3.2 Net Forces

Forces have magnitude and direction. The net force on an object is the sum of all the forces acting on the object. Objects change their speed and/or direction only when a net force is applied. If the net force on an object is zero, there is no change in motion (Newton's First Law).

P3.2 Net Forces

Forces have magnitude and direction. The net force on an object is the sum of all the forces acting on the object. Objects change their speed and/or direction only when a net force is applied. If the net force on an object is zero, there is no change in motion (Newton's First Law).

P3.2 Net Forces

Forces have magnitude and direction. The net force on an object is the sum of all the forces acting on the object. Objects change their speed and/or direction only when a net force is applied. If the net force on an object is zero, there is no change in motion (Newton's First Law).

P3.2 Net Forces

Forces have magnitude and direction. The net force on an object is the sum of all the forces acting on the object. Objects change their speed and/or direction only when a net force is applied. If the net force on an object is zero, there is no change in motion (Newton's First Law).

P3.2 Net Forces

Forces have magnitude and direction. The net force on an object is the sum of all the forces acting on the object. Objects change their speed and/or direction only when a net force is applied. If the net force on an object is zero, there is no change in motion (Newton's First Law).

P3.2 Net Forces

Forces have magnitude and direction. The net force on an object is the sum of all the forces acting on the object. Objects change their speed and/or direction only when a net force is applied. If the net force on an object is zero, there is no change in motion (Newton's First Law).



P3.2B Compare work done in different situations.



P3.2B Compare work done in different situations.



P3.2B Compare work done in different situations.



P3.2B Compare work done in different situations.



P3.2B Compare work done in different situations.



P3.2B Compare work done in different situations.

P3.2 Net Forces

Forces have magnitude and direction. The net force on an object is the sum of all the forces acting on the object. Objects change their speed and/or direction only when a net force is applied. If the net force on an object is zero, there is no change in motion (Newton's First Law).

P3.2 Net Forces

Forces have magnitude and direction. The net force on an object is the sum of all the forces acting on the object. Objects change their speed and/or direction only when a net force is applied. If the net force on an object is zero, there is no change in motion (Newton's First Law).

P3.2 Net Forces

Forces have magnitude and direction. The net force on an object is the sum of all the forces acting on the object. Objects change their speed and/or direction only when a net force is applied. If the net force on an object is zero, there is no change in motion (Newton's First Law).

P3.2 Net Forces

Forces have magnitude and direction. The net force on an object is the sum of all the forces acting on the object. Objects change their speed and/or direction only when a net force is applied. If the net force on an object is zero, there is no change in motion (Newton's First Law).

P3.2 Net Forces

Forces have magnitude and direction. The net force on an object is the sum of all the forces acting on the object. Objects change their speed and/or direction only when a net force is applied. If the net force on an object is zero, there is no change in motion (Newton's First Law).

P3.2 Net Forces

Forces have magnitude and direction. The net force on an object is the sum of all the forces acting on the object. Objects change their speed and/or direction only when a net force is applied. If the net force on an object is zero, there is no change in motion (Newton's First Law).



P3.2C Calculate the net force acting on an object.



P3.2C Calculate the net force acting on an object.



P3.2C Calculate the net force acting on an object.



P3.2C Calculate the net force acting on an object.



P3.2C Calculate the net force acting on an object.



P3.2C Calculate the net force acting on an object.

P3.2 Net Forces

Forces have magnitude and direction. The net force on an object is the sum of all the forces acting on the object. Objects change their speed and/or direction only when a net force is applied. If the net force on an object is zero, there is no change in motion (Newton's First Law).

P3.2 Net Forces

Forces have magnitude and direction. The net force on an object is the sum of all the forces acting on the object. Objects change their speed and/or direction only when a net force is applied. If the net force on an object is zero, there is no change in motion (Newton's First Law).

P3.2 Net Forces

Forces have magnitude and direction. The net force on an object is the sum of all the forces acting on the object. Objects change their speed and/or direction only when a net force is applied. If the net force on an object is zero, there is no change in motion (Newton's First Law).

P3.2 Net Forces

Forces have magnitude and direction. The net force on an object is the sum of all the forces acting on the object. Objects change their speed and/or direction only when a net force is applied. If the net force on an object is zero, there is no change in motion (Newton's First Law).

P3.2 Net Forces

Forces have magnitude and direction. The net force on an object is the sum of all the forces acting on the object. Objects change their speed and/or direction only when a net force is applied. If the net force on an object is zero, there is no change in motion (Newton's First Law).

P3.2 Net Forces

Forces have magnitude and direction. The net force on an object is the sum of all the forces acting on the object. Objects change their speed and/or direction only when a net force is applied. If the net force on an object is zero, there is no change in motion (Newton's First Law).



P3.2d Calculate all the forces on an object on an inclined plane and describe the object's motion based on the forces using free-body diagrams.



P3.2d Calculate all the forces on an object on an inclined plane and describe the object's motion based on the forces using free-body diagrams.



P3.2d Calculate all the forces on an object on an inclined plane and describe the object's motion based on the forces using free-body diagrams.



P3.2d Calculate all the forces on an object on an inclined plane and describe the object's motion based on the forces using free-body diagrams.



P3.2d Calculate all the forces on an object on an inclined plane and describe the object's motion based on the forces using free-body diagrams.



P3.2d Calculate all the forces on an object on an inclined plane and describe the object's motion based on the forces using free-body diagrams.

P3.3 Newton's Third Law

Whenever one object exerts a force on another object, a force equal in magnitude and opposite in direction is exerted back on the first object.

P3.3 Newton's Third Law

Whenever one object exerts a force on another object, a force equal in magnitude and opposite in direction is exerted back on the first object.

P3.3 Newton's Third Law

Whenever one object exerts a force on another object, a force equal in magnitude and opposite in direction is exerted back on the first object.

P3.3 Newton's Third Law

Whenever one object exerts a force on another object, a force equal in magnitude and opposite in direction is exerted back on the first object.

P3.3 Newton's Third Law

Whenever one object exerts a force on another object, a force equal in magnitude and opposite in direction is exerted back on the first object.

P3.3 Newton's Third Law

Whenever one object exerts a force on another object, a force equal in magnitude and opposite in direction is exerted back on the first object.



P3.3A Identify the action and reaction force from examples of forces in everyday situations (e.g., book on a table, walking across the floor, pushing open a door).



P3.3A Identify the action and reaction force from examples of forces in everyday situations (e.g., book on a table, walking across the floor, pushing open a door).



P3.3A Identify the action and reaction force from examples of forces in everyday situations (e.g., book on a table, walking across the floor, pushing open a door).



P3.3A Identify the action and reaction force from examples of forces in everyday situations (e.g., book on a table, walking across the floor, pushing open a door).



P3.3A Identify the action and reaction force from examples of forces in everyday situations (e.g., book on a table, walking across the floor, pushing open a door).



P3.3A Identify the action and reaction force from examples of forces in everyday situations (e.g., book on a table, walking across the floor, pushing open a door).

P3.3 Newton's Third Law

Whenever one object exerts a force on another object, a force equal in magnitude and opposite in direction is exerted back on the first object.

P3.3 Newton's Third Law

Whenever one object exerts a force on another object, a force equal in magnitude and opposite in direction is exerted back on the first object.

P3.3 Newton's Third Law

Whenever one object exerts a force on another object, a force equal in magnitude and opposite in direction is exerted back on the first object.

P3.3 Newton's Third Law

Whenever one object exerts a force on another object, a force equal in magnitude and opposite in direction is exerted back on the first object.

P3.3 Newton's Third Law

Whenever one object exerts a force on another object, a force equal in magnitude and opposite in direction is exerted back on the first object.

P3.3 Newton's Third Law

Whenever one object exerts a force on another object, a force equal in magnitude and opposite in direction is exerted back on the first object.



P3.3b Predict how the change in velocity of a small mass compares to the change in velocity of a large mass when the objects interact (e.g., collide).



P3.3b Predict how the change in velocity of a small mass compares to the change in velocity of a large mass when the objects interact (e.g., collide).



P3.3b Predict how the change in velocity of a small mass compares to the change in velocity of a large mass when the objects interact (e.g., collide).



P3.3b Predict how the change in velocity of a small mass compares to the change in velocity of a large mass when the objects interact (e.g., collide).



P3.3b Predict how the change in velocity of a small mass compares to the change in velocity of a large mass when the objects interact (e.g., collide).



P3.3b Predict how the change in velocity of a small mass compares to the change in velocity of a large mass when the objects interact (e.g., collide).

P3.3 Newton's Third Law

Whenever one object exerts a force on another object, a force equal in magnitude and opposite in direction is exerted back on the first object.

P3.3 Newton's Third Law

Whenever one object exerts a force on another object, a force equal in magnitude and opposite in direction is exerted back on the first object.

P3.3 Newton's Third Law

Whenever one object exerts a force on another object, a force equal in magnitude and opposite in direction is exerted back on the first object.

P3.3 Newton's Third Law

Whenever one object exerts a force on another object, a force equal in magnitude and opposite in direction is exerted back on the first object.

P3.3 Newton's Third Law

Whenever one object exerts a force on another object, a force equal in magnitude and opposite in direction is exerted back on the first object.

P3.3 Newton's Third Law

Whenever one object exerts a force on another object, a force equal in magnitude and opposite in direction is exerted back on the first object.



P3.3c Explain the recoil of a projectile launcher in terms of forces and masses.



P3.3c Explain the recoil of a projectile launcher in terms of forces and masses.



P3.3c Explain the recoil of a projectile launcher in terms of forces and masses.



P3.3c Explain the recoil of a projectile launcher in terms of forces and masses.



P3.3c Explain the recoil of a projectile launcher in terms of forces and masses.



P3.3c Explain the recoil of a projectile launcher in terms of forces and masses.

P3.3 Newton's Third Law

Whenever one object exerts a force on another object, a force equal in magnitude and opposite in direction is exerted back on the first object.

P3.3 Newton's Third Law

Whenever one object exerts a force on another object, a force equal in magnitude and opposite in direction is exerted back on the first object.

P3.3 Newton's Third Law

Whenever one object exerts a force on another object, a force equal in magnitude and opposite in direction is exerted back on the first object.

P3.3 Newton's Third Law

Whenever one object exerts a force on another object, a force equal in magnitude and opposite in direction is exerted back on the first object.

P3.3 Newton's Third Law

Whenever one object exerts a force on another object, a force equal in magnitude and opposite in direction is exerted back on the first object.

P3.3 Newton's Third Law

Whenever one object exerts a force on another object, a force equal in magnitude and opposite in direction is exerted back on the first object.



P3.3d Analyze why seat belts may be more important in autos than in buses.



P3.3d Analyze why seat belts may be more important in autos than in buses.



P3.3d Analyze why seat belts may be more important in autos than in buses.



P3.3d Analyze why seat belts may be more important in autos than in buses.



P3.3d Analyze why seat belts may be more important in autos than in buses.



P3.3d Analyze why seat belts may be more important in autos than in buses.

P3.4 Forces and Acceleration

The change of speed and/or direction (acceleration) of an object is proportional to the net force and inversely proportional to the mass of the object. The acceleration and net force are always in the same direction.

P3.4 Forces and Acceleration

The change of speed and/or direction (acceleration) of an object is proportional to the net force and inversely proportional to the mass of the object. The acceleration and net force are always in the same direction.

P3.4 Forces and Acceleration

The change of speed and/or direction (acceleration) of an object is proportional to the net force and inversely proportional to the mass of the object. The acceleration and net force are always in the same direction.

P3.4 Forces and Acceleration

The change of speed and/or direction (acceleration) of an object is proportional to the net force and inversely proportional to the mass of the object. The acceleration and net force are always in the same direction.

P3.4 Forces and Acceleration

The change of speed and/or direction (acceleration) of an object is proportional to the net force and inversely proportional to the mass of the object. The acceleration and net force are always in the same direction.

P3.4 Forces and Acceleration

The change of speed and/or direction (acceleration) of an object is proportional to the net force and inversely proportional to the mass of the object. The acceleration and net force are always in the same direction.

E

P3.4A Predict the change in motion of an object acted on by several forces.

E

P3.4A Predict the change in motion of an object acted on by several forces.

E

P3.4A Predict the change in motion of an object acted on by several forces.

E

P3.4A Predict the change in motion of an object acted on by several forces.

E

P3.4A Predict the change in motion of an object acted on by several forces.

E

P3.4A Predict the change in motion of an object acted on by several forces.

P3.4 Forces and Acceleration

The change of speed and/or direction (acceleration) of an object is proportional to the net force and inversely proportional to the mass of the object. The acceleration and net force are always in the same direction.

P3.4 Forces and Acceleration

The change of speed and/or direction (acceleration) of an object is proportional to the net force and inversely proportional to the mass of the object. The acceleration and net force are always in the same direction.

P3.4 Forces and Acceleration

The change of speed and/or direction (acceleration) of an object is proportional to the net force and inversely proportional to the mass of the object. The acceleration and net force are always in the same direction.

P3.4 Forces and Acceleration

The change of speed and/or direction (acceleration) of an object is proportional to the net force and inversely proportional to the mass of the object. The acceleration and net force are always in the same direction.

P3.4 Forces and Acceleration

The change of speed and/or direction (acceleration) of an object is proportional to the net force and inversely proportional to the mass of the object. The acceleration and net force are always in the same direction.

P3.4 Forces and Acceleration

The change of speed and/or direction (acceleration) of an object is proportional to the net force and inversely proportional to the mass of the object. The acceleration and net force are always in the same direction.



P3.4B Identify forces acting on objects moving with constant velocity (e.g., cars on a highway).



P3.4B Identify forces acting on objects moving with constant velocity (e.g., cars on a highway).



P3.4B Identify forces acting on objects moving with constant velocity (e.g., cars on a highway).



P3.4B Identify forces acting on objects moving with constant velocity (e.g., cars on a highway).



P3.4B Identify forces acting on objects moving with constant velocity (e.g., cars on a highway).



P3.4B Identify forces acting on objects moving with constant velocity (e.g., cars on a highway).

P3.4 Forces and Acceleration

The change of speed and/or direction (acceleration) of an object is proportional to the net force and inversely proportional to the mass of the object. The acceleration and net force are always in the same direction.

P3.4 Forces and Acceleration

The change of speed and/or direction (acceleration) of an object is proportional to the net force and inversely proportional to the mass of the object. The acceleration and net force are always in the same direction.

P3.4 Forces and Acceleration

The change of speed and/or direction (acceleration) of an object is proportional to the net force and inversely proportional to the mass of the object. The acceleration and net force are always in the same direction.

P3.4 Forces and Acceleration

The change of speed and/or direction (acceleration) of an object is proportional to the net force and inversely proportional to the mass of the object. The acceleration and net force are always in the same direction.

P3.4 Forces and Acceleration

The change of speed and/or direction (acceleration) of an object is proportional to the net force and inversely proportional to the mass of the object. The acceleration and net force are always in the same direction.

P3.4 Forces and Acceleration

The change of speed and/or direction (acceleration) of an object is proportional to the net force and inversely proportional to the mass of the object. The acceleration and net force are always in the same direction.

E

P3.4C Solve problems involving force, mass, and acceleration in linear motion (Newton's second law).

E

P3.4C Solve problems involving force, mass, and acceleration in linear motion (Newton's second law).

E

P3.4C Solve problems involving force, mass, and acceleration in linear motion (Newton's second law).

E

P3.4C Solve problems involving force, mass, and acceleration in linear motion (Newton's second law).

E

P3.4C Solve problems involving force, mass, and acceleration in linear motion (Newton's second law).

E

P3.4C Solve problems involving force, mass, and acceleration in linear motion (Newton's second law).

P3.4 Forces and Acceleration

The change of speed and/or direction (acceleration) of an object is proportional to the net force and inversely proportional to the mass of the object. The acceleration and net force are always in the same direction.

P3.4 Forces and Acceleration

The change of speed and/or direction (acceleration) of an object is proportional to the net force and inversely proportional to the mass of the object. The acceleration and net force are always in the same direction.

P3.4 Forces and Acceleration

The change of speed and/or direction (acceleration) of an object is proportional to the net force and inversely proportional to the mass of the object. The acceleration and net force are always in the same direction.

P3.4 Forces and Acceleration

The change of speed and/or direction (acceleration) of an object is proportional to the net force and inversely proportional to the mass of the object. The acceleration and net force are always in the same direction.

P3.4 Forces and Acceleration

The change of speed and/or direction (acceleration) of an object is proportional to the net force and inversely proportional to the mass of the object. The acceleration and net force are always in the same direction.

P3.4 Forces and Acceleration

The change of speed and/or direction (acceleration) of an object is proportional to the net force and inversely proportional to the mass of the object. The acceleration and net force are always in the same direction.



P3.4D Identify the force(s) acting on objects moving with uniform circular motion (e.g., a car on a circular track, satellites in orbit).



P3.4D Identify the force(s) acting on objects moving with uniform circular motion (e.g., a car on a circular track, satellites in orbit).



P3.4D Identify the force(s) acting on objects moving with uniform circular motion (e.g., a car on a circular track, satellites in orbit).



P3.4D Identify the force(s) acting on objects moving with uniform circular motion (e.g., a car on a circular track, satellites in orbit).



P3.4D Identify the force(s) acting on objects moving with uniform circular motion (e.g., a car on a circular track, satellites in orbit).



P3.4D Identify the force(s) acting on objects moving with uniform circular motion (e.g., a car on a circular track, satellites in orbit).

P3.4 Forces and Acceleration

The change of speed and/or direction (acceleration) of an object is proportional to the net force and inversely proportional to the mass of the object. The acceleration and net force are always in the same direction.

P3.4 Forces and Acceleration

The change of speed and/or direction (acceleration) of an object is proportional to the net force and inversely proportional to the mass of the object. The acceleration and net force are always in the same direction.

P3.4 Forces and Acceleration

The change of speed and/or direction (acceleration) of an object is proportional to the net force and inversely proportional to the mass of the object. The acceleration and net force are always in the same direction.

P3.4 Forces and Acceleration

The change of speed and/or direction (acceleration) of an object is proportional to the net force and inversely proportional to the mass of the object. The acceleration and net force are always in the same direction.

P3.4 Forces and Acceleration

The change of speed and/or direction (acceleration) of an object is proportional to the net force and inversely proportional to the mass of the object. The acceleration and net force are always in the same direction.

P3.4 Forces and Acceleration

The change of speed and/or direction (acceleration) of an object is proportional to the net force and inversely proportional to the mass of the object. The acceleration and net force are always in the same direction.



P3.4e Solve problems involving force, mass, and acceleration in two-dimensional projectile motion restricted to an initial horizontal velocity with no initial vertical velocity (e.g., ball rolling off a table).



P3.4e Solve problems involving force, mass, and acceleration in two-dimensional projectile motion restricted to an initial horizontal velocity with no initial vertical velocity (e.g., ball rolling off a table).



P3.4e Solve problems involving force, mass, and acceleration in two-dimensional projectile motion restricted to an initial horizontal velocity with no initial vertical velocity (e.g., ball rolling off a table).



P3.4e Solve problems involving force, mass, and acceleration in two-dimensional projectile motion restricted to an initial horizontal velocity with no initial vertical velocity (e.g., ball rolling off a table).



P3.4e Solve problems involving force, mass, and acceleration in two-dimensional projectile motion restricted to an initial horizontal velocity with no initial vertical velocity (e.g., ball rolling off a table).



P3.4e Solve problems involving force, mass, and acceleration in two-dimensional projectile motion restricted to an initial horizontal velocity with no initial vertical velocity (e.g., ball rolling off a table).

P3.4 Forces and Acceleration

The change of speed and/or direction (acceleration) of an object is proportional to the net force and inversely proportional to the mass of the object. The acceleration and net force are always in the same direction.

P3.4 Forces and Acceleration

The change of speed and/or direction (acceleration) of an object is proportional to the net force and inversely proportional to the mass of the object. The acceleration and net force are always in the same direction.

P3.4 Forces and Acceleration

The change of speed and/or direction (acceleration) of an object is proportional to the net force and inversely proportional to the mass of the object. The acceleration and net force are always in the same direction.

P3.4 Forces and Acceleration

The change of speed and/or direction (acceleration) of an object is proportional to the net force and inversely proportional to the mass of the object. The acceleration and net force are always in the same direction.

P3.4 Forces and Acceleration

The change of speed and/or direction (acceleration) of an object is proportional to the net force and inversely proportional to the mass of the object. The acceleration and net force are always in the same direction.

P3.4 Forces and Acceleration

The change of speed and/or direction (acceleration) of an object is proportional to the net force and inversely proportional to the mass of the object. The acceleration and net force are always in the same direction.



P3.4f Calculate the changes in velocity of a thrown or hit object during and after the time it is acted on by the force.



P3.4f Calculate the changes in velocity of a thrown or hit object during and after the time it is acted on by the force.



P3.4f Calculate the changes in velocity of a thrown or hit object during and after the time it is acted on by the force.



P3.4f Calculate the changes in velocity of a thrown or hit object during and after the time it is acted on by the force.



P3.4f Calculate the changes in velocity of a thrown or hit object during and after the time it is acted on by the force.



P3.4f Calculate the changes in velocity of a thrown or hit object during and after the time it is acted on by the force.

P3.4 Forces and Acceleration

The change of speed and/or direction (acceleration) of an object is proportional to the net force and inversely proportional to the mass of the object. The acceleration and net force are always in the same direction.

P3.4 Forces and Acceleration

The change of speed and/or direction (acceleration) of an object is proportional to the net force and inversely proportional to the mass of the object. The acceleration and net force are always in the same direction.

P3.4 Forces and Acceleration

The change of speed and/or direction (acceleration) of an object is proportional to the net force and inversely proportional to the mass of the object. The acceleration and net force are always in the same direction.

P3.4 Forces and Acceleration

The change of speed and/or direction (acceleration) of an object is proportional to the net force and inversely proportional to the mass of the object. The acceleration and net force are always in the same direction.

P3.4 Forces and Acceleration

The change of speed and/or direction (acceleration) of an object is proportional to the net force and inversely proportional to the mass of the object. The acceleration and net force are always in the same direction.

P3.4 Forces and Acceleration

The change of speed and/or direction (acceleration) of an object is proportional to the net force and inversely proportional to the mass of the object. The acceleration and net force are always in the same direction.



P3.4g Explain how the time of impact can affect the net force (e.g., air bags in cars, catching a ball).



P3.4g Explain how the time of impact can affect the net force (e.g., air bags in cars, catching a ball).



P3.4g Explain how the time of impact can affect the net force (e.g., air bags in cars, catching a ball).



P3.4g Explain how the time of impact can affect the net force (e.g., air bags in cars, catching a ball).



P3.4g Explain how the time of impact can affect the net force (e.g., air bags in cars, catching a ball).



P3.4g Explain how the time of impact can affect the net force (e.g., air bags in cars, catching a ball).

P3.5x Momentum

A moving object has a quantity of motion (momentum) that depends on its velocity and mass. In interactions between objects, the total momentum of the objects does not change.

P3.5x Momentum

A moving object has a quantity of motion (momentum) that depends on its velocity and mass. In interactions between objects, the total momentum of the objects does not change.

P3.5x Momentum

A moving object has a quantity of motion (momentum) that depends on its velocity and mass. In interactions between objects, the total momentum of the objects does not change.

P3.5x Momentum

A moving object has a quantity of motion (momentum) that depends on its velocity and mass. In interactions between objects, the total momentum of the objects does not change.

P3.5x Momentum

A moving object has a quantity of motion (momentum) that depends on its velocity and mass. In interactions between objects, the total momentum of the objects does not change.

P3.5x Momentum

A moving object has a quantity of motion (momentum) that depends on its velocity and mass. In interactions between objects, the total momentum of the objects does not change.



P3.5a Apply conservation of momentum to solve simple collision problems.



P3.5a Apply conservation of momentum to solve simple collision problems.



P3.5a Apply conservation of momentum to solve simple collision problems.



P3.5a Apply conservation of momentum to solve simple collision problems.



P3.5a Apply conservation of momentum to solve simple collision problems.



P3.5a Apply conservation of momentum to solve simple collision problems.

P3.6 Gravitational Interactions

Gravitation is a universal attractive force that a mass exerts on every other mass. The strength of the gravitational force between two masses is proportional to the masses and inversely proportional to the square of the distance between them.

P3.6 Gravitational Interactions

Gravitation is a universal attractive force that a mass exerts on every other mass. The strength of the gravitational force between two masses is proportional to the masses and inversely proportional to the square of the distance between them.

P3.6 Gravitational Interactions

Gravitation is a universal attractive force that a mass exerts on every other mass. The strength of the gravitational force between two masses is proportional to the masses and inversely proportional to the square of the distance between them.

P3.6 Gravitational Interactions

Gravitation is a universal attractive force that a mass exerts on every other mass. The strength of the gravitational force between two masses is proportional to the masses and inversely proportional to the square of the distance between them.

P3.6 Gravitational Interactions

Gravitation is a universal attractive force that a mass exerts on every other mass. The strength of the gravitational force between two masses is proportional to the masses and inversely proportional to the square of the distance between them.

P3.6 Gravitational Interactions

Gravitation is a universal attractive force that a mass exerts on every other mass. The strength of the gravitational force between two masses is proportional to the masses and inversely proportional to the square of the distance between them.

E

P3.6A Explain earth-moon interactions (orbital motion) in terms of forces.

E

P3.6A Explain earth-moon interactions (orbital motion) in terms of forces.

E

P3.6A Explain earth-moon interactions (orbital motion) in terms of forces.

E

P3.6A Explain earth-moon interactions (orbital motion) in terms of forces.

E

P3.6A Explain earth-moon interactions (orbital motion) in terms of forces.

E

P3.6A Explain earth-moon interactions (orbital motion) in terms of forces.

P3.6 Gravitational Interactions

Gravitation is a universal attractive force that a mass exerts on every other mass. The strength of the gravitational force between two masses is proportional to the masses and inversely proportional to the square of the distance between them.

P3.6 Gravitational Interactions

Gravitation is a universal attractive force that a mass exerts on every other mass. The strength of the gravitational force between two masses is proportional to the masses and inversely proportional to the square of the distance between them.

P3.6 Gravitational Interactions

Gravitation is a universal attractive force that a mass exerts on every other mass. The strength of the gravitational force between two masses is proportional to the masses and inversely proportional to the square of the distance between them.

P3.6 Gravitational Interactions

Gravitation is a universal attractive force that a mass exerts on every other mass. The strength of the gravitational force between two masses is proportional to the masses and inversely proportional to the square of the distance between them.

P3.6 Gravitational Interactions

Gravitation is a universal attractive force that a mass exerts on every other mass. The strength of the gravitational force between two masses is proportional to the masses and inversely proportional to the square of the distance between them.

P3.6 Gravitational Interactions

Gravitation is a universal attractive force that a mass exerts on every other mass. The strength of the gravitational force between two masses is proportional to the masses and inversely proportional to the square of the distance between them.



P3.6B Predict how the gravitational force between objects changes when the distance between them changes.



P3.6B Predict how the gravitational force between objects changes when the distance between them changes.



P3.6B Predict how the gravitational force between objects changes when the distance between them changes.



P3.6B Predict how the gravitational force between objects changes when the distance between them changes.



P3.6B Predict how the gravitational force between objects changes when the distance between them changes.



P3.6B Predict how the gravitational force between objects changes when the distance between them changes.

P3.6 Gravitational Interactions

Gravitation is a universal attractive force that a mass exerts on every other mass. The strength of the gravitational force between two masses is proportional to the masses and inversely proportional to the square of the distance between them.

P3.6 Gravitational Interactions

Gravitation is a universal attractive force that a mass exerts on every other mass. The strength of the gravitational force between two masses is proportional to the masses and inversely proportional to the square of the distance between them.

P3.6 Gravitational Interactions

Gravitation is a universal attractive force that a mass exerts on every other mass. The strength of the gravitational force between two masses is proportional to the masses and inversely proportional to the square of the distance between them.

P3.6 Gravitational Interactions

Gravitation is a universal attractive force that a mass exerts on every other mass. The strength of the gravitational force between two masses is proportional to the masses and inversely proportional to the square of the distance between them.

P3.6 Gravitational Interactions

Gravitation is a universal attractive force that a mass exerts on every other mass. The strength of the gravitational force between two masses is proportional to the masses and inversely proportional to the square of the distance between them.

P3.6 Gravitational Interactions

Gravitation is a universal attractive force that a mass exerts on every other mass. The strength of the gravitational force between two masses is proportional to the masses and inversely proportional to the square of the distance between them.



P3.6C Explain how your weight on Earth could be different from your weight on another planet.



P3.6C Explain how your weight on Earth could be different from your weight on another planet.



P3.6C Explain how your weight on Earth could be different from your weight on another planet.



P3.6C Explain how your weight on Earth could be different from your weight on another planet.



P3.6C Explain how your weight on Earth could be different from your weight on another planet.



P3.6C Explain how your weight on Earth could be different from your weight on another planet.

P3.6 Gravitational Interactions

Gravitation is a universal attractive force that a mass exerts on every other mass. The strength of the gravitational force between two masses is proportional to the masses and inversely proportional to the square of the distance between them.

P3.6 Gravitational Interactions

Gravitation is a universal attractive force that a mass exerts on every other mass. The strength of the gravitational force between two masses is proportional to the masses and inversely proportional to the square of the distance between them.

P3.6 Gravitational Interactions

Gravitation is a universal attractive force that a mass exerts on every other mass. The strength of the gravitational force between two masses is proportional to the masses and inversely proportional to the square of the distance between them.

P3.6 Gravitational Interactions

Gravitation is a universal attractive force that a mass exerts on every other mass. The strength of the gravitational force between two masses is proportional to the masses and inversely proportional to the square of the distance between them.

P3.6 Gravitational Interactions

Gravitation is a universal attractive force that a mass exerts on every other mass. The strength of the gravitational force between two masses is proportional to the masses and inversely proportional to the square of the distance between them.

P3.6 Gravitational Interactions

Gravitation is a universal attractive force that a mass exerts on every other mass. The strength of the gravitational force between two masses is proportional to the masses and inversely proportional to the square of the distance between them.



P3.6d Calculate force, masses, or distance, given any three of these quantities, by applying the Law of Universal Gravitation, given the value of G .



P3.6d Calculate force, masses, or distance, given any three of these quantities, by applying the Law of Universal Gravitation, given the value of G .



P3.6d Calculate force, masses, or distance, given any three of these quantities, by applying the Law of Universal Gravitation, given the value of G .



P3.6d Calculate force, masses, or distance, given any three of these quantities, by applying the Law of Universal Gravitation, given the value of G .



P3.6d Calculate force, masses, or distance, given any three of these quantities, by applying the Law of Universal Gravitation, given the value of G .



P3.6d Calculate force, masses, or distance, given any three of these quantities, by applying the Law of Universal Gravitation, given the value of G .

P3.6 Gravitational Interactions

Gravitation is a universal attractive force that a mass exerts on every other mass. The strength of the gravitational force between two masses is proportional to the masses and inversely proportional to the square of the distance between them.

P3.6 Gravitational Interactions

Gravitation is a universal attractive force that a mass exerts on every other mass. The strength of the gravitational force between two masses is proportional to the masses and inversely proportional to the square of the distance between them.

P3.6 Gravitational Interactions

Gravitation is a universal attractive force that a mass exerts on every other mass. The strength of the gravitational force between two masses is proportional to the masses and inversely proportional to the square of the distance between them.

P3.6 Gravitational Interactions

Gravitation is a universal attractive force that a mass exerts on every other mass. The strength of the gravitational force between two masses is proportional to the masses and inversely proportional to the square of the distance between them.

P3.6 Gravitational Interactions

Gravitation is a universal attractive force that a mass exerts on every other mass. The strength of the gravitational force between two masses is proportional to the masses and inversely proportional to the square of the distance between them.

P3.6 Gravitational Interactions

Gravitation is a universal attractive force that a mass exerts on every other mass. The strength of the gravitational force between two masses is proportional to the masses and inversely proportional to the square of the distance between them.



P3.6e Draw arrows (vectors) to represent how the direction and magnitude of a force changes on an object in an elliptical orbit.



P3.6e Draw arrows (vectors) to represent how the direction and magnitude of a force changes on an object in an elliptical orbit.



P3.6e Draw arrows (vectors) to represent how the direction and magnitude of a force changes on an object in an elliptical orbit.



P3.6e Draw arrows (vectors) to represent how the direction and magnitude of a force changes on an object in an elliptical orbit.



P3.6e Draw arrows (vectors) to represent how the direction and magnitude of a force changes on an object in an elliptical orbit.



P3.6e Draw arrows (vectors) to represent how the direction and magnitude of a force changes on an object in an elliptical orbit.

P3.7 Electric Charges

Electric force exists between any two charged objects. Oppositely charged objects attract, while objects with like charge repel. The strength of the electric force between two charged objects is proportional to the magnitudes of the charges and inversely proportional to the square of the distance between them (Coulomb's Law).

P3.7 Electric Charges

Electric force exists between any two charged objects. Oppositely charged objects attract, while objects with like charge repel. The strength of the electric force between two charged objects is proportional to the magnitudes of the charges and inversely proportional to the square of the distance between them (Coulomb's Law).

P3.7 Electric Charges

Electric force exists between any two charged objects. Oppositely charged objects attract, while objects with like charge repel. The strength of the electric force between two charged objects is proportional to the magnitudes of the charges and inversely proportional to the square of the distance between them (Coulomb's Law).

P3.7 Electric Charges

Electric force exists between any two charged objects. Oppositely charged objects attract, while objects with like charge repel. The strength of the electric force between two charged objects is proportional to the magnitudes of the charges and inversely proportional to the square of the distance between them (Coulomb's Law).

P3.7 Electric Charges

Electric force exists between any two charged objects. Oppositely charged objects attract, while objects with like charge repel. The strength of the electric force between two charged objects is proportional to the magnitudes of the charges and inversely proportional to the square of the distance between them (Coulomb's Law).

P3.7 Electric Charges

Electric force exists between any two charged objects. Oppositely charged objects attract, while objects with like charge repel. The strength of the electric force between two charged objects is proportional to the magnitudes of the charges and inversely proportional to the square of the distance between them (Coulomb's Law).



P3.7A Predict how the electric force between charged objects varies when the distance between them and/or the magnitude of charges change.



P3.7A Predict how the electric force between charged objects varies when the distance between them and/or the magnitude of charges change.



P3.7A Predict how the electric force between charged objects varies when the distance between them and/or the magnitude of charges change.



P3.7A Predict how the electric force between charged objects varies when the distance between them and/or the magnitude of charges change.



P3.7A Predict how the electric force between charged objects varies when the distance between them and/or the magnitude of charges change.



P3.7A Predict how the electric force between charged objects varies when the distance between them and/or the magnitude of charges change.

P3.7 Electric Charges

Electric force exists between any two charged objects. Oppositely charged objects attract, while objects with like charge repel. The strength of the electric force between two charged objects is proportional to the magnitudes of the charges and inversely proportional to the square of the distance between them (Coulomb's Law).

P3.7 Electric Charges

Electric force exists between any two charged objects. Oppositely charged objects attract, while objects with like charge repel. The strength of the electric force between two charged objects is proportional to the magnitudes of the charges and inversely proportional to the square of the distance between them (Coulomb's Law).

P3.7 Electric Charges

Electric force exists between any two charged objects. Oppositely charged objects attract, while objects with like charge repel. The strength of the electric force between two charged objects is proportional to the magnitudes of the charges and inversely proportional to the square of the distance between them (Coulomb's Law).

P3.7 Electric Charges

Electric force exists between any two charged objects. Oppositely charged objects attract, while objects with like charge repel. The strength of the electric force between two charged objects is proportional to the magnitudes of the charges and inversely proportional to the square of the distance between them (Coulomb's Law).

P3.7 Electric Charges

Electric force exists between any two charged objects. Oppositely charged objects attract, while objects with like charge repel. The strength of the electric force between two charged objects is proportional to the magnitudes of the charges and inversely proportional to the square of the distance between them (Coulomb's Law).

P3.7 Electric Charges

Electric force exists between any two charged objects. Oppositely charged objects attract, while objects with like charge repel. The strength of the electric force between two charged objects is proportional to the magnitudes of the charges and inversely proportional to the square of the distance between them (Coulomb's Law).

E

P3.7B Explain why acquiring a large excess static charge (e.g., pulling off a wool cap, touching a Van de Graaff generator, combing) affects your hair.

E

P3.7B Explain why acquiring a large excess static charge (e.g., pulling off a wool cap, touching a Van de Graaff generator, combing) affects your hair.

E

P3.7B Explain why acquiring a large excess static charge (e.g., pulling off a wool cap, touching a Van de Graaff generator, combing) affects your hair.

E

P3.7B Explain why acquiring a large excess static charge (e.g., pulling off a wool cap, touching a Van de Graaff generator, combing) affects your hair.

E

P3.7B Explain why acquiring a large excess static charge (e.g., pulling off a wool cap, touching a Van de Graaff generator, combing) affects your hair.

E

P3.7B Explain why acquiring a large excess static charge (e.g., pulling off a wool cap, touching a Van de Graaff generator, combing) affects your hair.

P3.7x Electric Charges — Interactions

Charged objects can attract electrically neutral objects by induction.

P3.7x Electric Charges — Interactions

Charged objects can attract electrically neutral objects by induction.

P3.7x Electric Charges — Interactions

Charged objects can attract electrically neutral objects by induction.

P3.7x Electric Charges — Interactions

Charged objects can attract electrically neutral objects by induction.

P3.7x Electric Charges — Interactions

Charged objects can attract electrically neutral objects by induction.

P3.7x Electric Charges — Interactions

Charged objects can attract electrically neutral objects by induction.



P3.7c Draw the redistribution of electric charges on a neutral object when a charged object is brought near.



P3.7c Draw the redistribution of electric charges on a neutral object when a charged object is brought near.



P3.7c Draw the redistribution of electric charges on a neutral object when a charged object is brought near.



P3.7c Draw the redistribution of electric charges on a neutral object when a charged object is brought near.



P3.7c Draw the redistribution of electric charges on a neutral object when a charged object is brought near.



P3.7c Draw the redistribution of electric charges on a neutral object when a charged object is brought near.

P3.7x Electric Charges — Interactions

Charged objects can attract electrically neutral objects by induction.

P3.7x Electric Charges — Interactions

Charged objects can attract electrically neutral objects by induction.

P3.7x Electric Charges — Interactions

Charged objects can attract electrically neutral objects by induction.

P3.7x Electric Charges — Interactions

Charged objects can attract electrically neutral objects by induction.

P3.7x Electric Charges — Interactions

Charged objects can attract electrically neutral objects by induction.

P3.7x Electric Charges — Interactions

Charged objects can attract electrically neutral objects by induction.



P3.7d Identify examples of induced static charges.



P3.7d Identify examples of induced static charges.



P3.7d Identify examples of induced static charges.



P3.7d Identify examples of induced static charges.



P3.7d Identify examples of induced static charges.



P3.7d Identify examples of induced static charges.

P3.7x Electric Charges — Interactions

Charged objects can attract electrically neutral objects by induction.

P3.7x Electric Charges — Interactions

Charged objects can attract electrically neutral objects by induction.

P3.7x Electric Charges — Interactions

Charged objects can attract electrically neutral objects by induction.

P3.7x Electric Charges — Interactions

Charged objects can attract electrically neutral objects by induction.

P3.7x Electric Charges — Interactions

Charged objects can attract electrically neutral objects by induction.

P3.7x Electric Charges — Interactions

Charged objects can attract electrically neutral objects by induction.



P3.7e Explain why an attractive force results from bringing a charged object near a neutral object.



P3.7e Explain why an attractive force results from bringing a charged object near a neutral object.



P3.7e Explain why an attractive force results from bringing a charged object near a neutral object.



P3.7e Explain why an attractive force results from bringing a charged object near a neutral object.



P3.7e Explain why an attractive force results from bringing a charged object near a neutral object.



P3.7e Explain why an attractive force results from bringing a charged object near a neutral object.

P3.7x Electric Charges — Interactions

Charged objects can attract electrically neutral objects by induction.

P3.7x Electric Charges — Interactions

Charged objects can attract electrically neutral objects by induction.

P3.7x Electric Charges — Interactions

Charged objects can attract electrically neutral objects by induction.

P3.7x Electric Charges — Interactions

Charged objects can attract electrically neutral objects by induction.

P3.7x Electric Charges — Interactions

Charged objects can attract electrically neutral objects by induction.

P3.7x Electric Charges — Interactions

Charged objects can attract electrically neutral objects by induction.



P3.7f Determine the new electric force on charged objects after they touch and are then separated.



P3.7f Determine the new electric force on charged objects after they touch and are then separated.



P3.7f Determine the new electric force on charged objects after they touch and are then separated.



P3.7f Determine the new electric force on charged objects after they touch and are then separated.



P3.7f Determine the new electric force on charged objects after they touch and are then separated.



P3.7f Determine the new electric force on charged objects after they touch and are then separated.

P3.7x Electric Charges — Interactions

Charged objects can attract electrically neutral objects by induction.

P3.7x Electric Charges — Interactions

Charged objects can attract electrically neutral objects by induction.

P3.7x Electric Charges — Interactions

Charged objects can attract electrically neutral objects by induction.

P3.7x Electric Charges — Interactions

Charged objects can attract electrically neutral objects by induction.

P3.7x Electric Charges — Interactions

Charged objects can attract electrically neutral objects by induction.

P3.7x Electric Charges — Interactions

Charged objects can attract electrically neutral objects by induction.



P3.7g Propose a mechanism based on electric forces to explain current flow in an electric circuit.



P3.7g Propose a mechanism based on electric forces to explain current flow in an electric circuit.



P3.7g Propose a mechanism based on electric forces to explain current flow in an electric circuit.



P3.7g Propose a mechanism based on electric forces to explain current flow in an electric circuit.



P3.7g Propose a mechanism based on electric forces to explain current flow in an electric circuit.



P3.7g Propose a mechanism based on electric forces to explain current flow in an electric circuit.

P3.p8 Magnetic Force (prerequisite)

Magnets exert forces on all objects made of ferromagnetic materials (e.g., iron, cobalt, and nickel) as well as other magnets. This force acts at a distance. Magnetic fields accompany magnets and are related to the strength and direction of the magnetic force. (prerequisite)

P3.p8 Magnetic Force (prerequisite)

Magnets exert forces on all objects made of ferromagnetic materials (e.g., iron, cobalt, and nickel) as well as other magnets. This force acts at a distance. Magnetic fields accompany magnets and are related to the strength and direction of the magnetic force. (prerequisite)

P3.p8 Magnetic Force (prerequisite)

Magnets exert forces on all objects made of ferromagnetic materials (e.g., iron, cobalt, and nickel) as well as other magnets. This force acts at a distance. Magnetic fields accompany magnets and are related to the strength and direction of the magnetic force. (prerequisite)

P3.p8 Magnetic Force (prerequisite)

Magnets exert forces on all objects made of ferromagnetic materials (e.g., iron, cobalt, and nickel) as well as other magnets. This force acts at a distance. Magnetic fields accompany magnets and are related to the strength and direction of the magnetic force. (prerequisite)

P3.p8 Magnetic Force (prerequisite)

Magnets exert forces on all objects made of ferromagnetic materials (e.g., iron, cobalt, and nickel) as well as other magnets. This force acts at a distance. Magnetic fields accompany magnets and are related to the strength and direction of the magnetic force. (prerequisite)

P3.p8 Magnetic Force (prerequisite)

Magnets exert forces on all objects made of ferromagnetic materials (e.g., iron, cobalt, and nickel) as well as other magnets. This force acts at a distance. Magnetic fields accompany magnets and are related to the strength and direction of the magnetic force. (prerequisite)



P3.p8A Create a representation of magnetic field lines around a bar magnet and qualitatively describe how the relative strength and direction of the magnetic force changes at various places in the field. (prerequisite)



P3.p8A Create a representation of magnetic field lines around a bar magnet and qualitatively describe how the relative strength and direction of the magnetic force changes at various places in the field. (prerequisite)



P3.p8A Create a representation of magnetic field lines around a bar magnet and qualitatively describe how the relative strength and direction of the magnetic force changes at various places in the field. (prerequisite)



P3.p8A Create a representation of magnetic field lines around a bar magnet and qualitatively describe how the relative strength and direction of the magnetic force changes at various places in the field. (prerequisite)



P3.p8A Create a representation of magnetic field lines around a bar magnet and qualitatively describe how the relative strength and direction of the magnetic force changes at various places in the field. (prerequisite)



P3.p8A Create a representation of magnetic field lines around a bar magnet and qualitatively describe how the relative strength and direction of the magnetic force changes at various places in the field. (prerequisite)

P3.8x Electromagnetic Force

Magnetic and electric forces are two aspects of a single electromagnetic force. Moving electric charges produce magnetic forces and moving magnets produce electric forces (e.g., electric current in a conductor).

P3.8x Electromagnetic Force

Magnetic and electric forces are two aspects of a single electromagnetic force. Moving electric charges produce magnetic forces and moving magnets produce electric forces (e.g., electric current in a conductor).

P3.8x Electromagnetic Force

Magnetic and electric forces are two aspects of a single electromagnetic force. Moving electric charges produce magnetic forces and moving magnets produce electric forces (e.g., electric current in a conductor).

P3.8x Electromagnetic Force

Magnetic and electric forces are two aspects of a single electromagnetic force. Moving electric charges produce magnetic forces and moving magnets produce electric forces (e.g., electric current in a conductor).

P3.8x Electromagnetic Force

Magnetic and electric forces are two aspects of a single electromagnetic force. Moving electric charges produce magnetic forces and moving magnets produce electric forces (e.g., electric current in a conductor).

P3.8x Electromagnetic Force

Magnetic and electric forces are two aspects of a single electromagnetic force. Moving electric charges produce magnetic forces and moving magnets produce electric forces (e.g., electric current in a conductor).



P3.8b Explain how the interaction of electric and magnetic forces is the basis for electric motors, generators, and the production of electromagnetic waves.



P3.8b Explain how the interaction of electric and magnetic forces is the basis for electric motors, generators, and the production of electromagnetic waves.



P3.8b Explain how the interaction of electric and magnetic forces is the basis for electric motors, generators, and the production of electromagnetic waves.



P3.8b Explain how the interaction of electric and magnetic forces is the basis for electric motors, generators, and the production of electromagnetic waves.



P3.8b Explain how the interaction of electric and magnetic forces is the basis for electric motors, generators, and the production of electromagnetic waves.



P3.8b Explain how the interaction of electric and magnetic forces is the basis for electric motors, generators, and the production of electromagnetic waves.

P4.1 Energy Transfer

Moving objects and waves transfer energy from one location to another. They also transfer energy to objects during interactions (e.g., sunlight transfers energy to the ground when it warms the ground; sunlight also transfers energy from the Sun to the Earth).

P4.1 Energy Transfer

Moving objects and waves transfer energy from one location to another. They also transfer energy to objects during interactions (e.g., sunlight transfers energy to the ground when it warms the ground; sunlight also transfers energy from the Sun to the Earth).

P4.1 Energy Transfer

Moving objects and waves transfer energy from one location to another. They also transfer energy to objects during interactions (e.g., sunlight transfers energy to the ground when it warms the ground; sunlight also transfers energy from the Sun to the Earth).

P4.1 Energy Transfer

Moving objects and waves transfer energy from one location to another. They also transfer energy to objects during interactions (e.g., sunlight transfers energy to the ground when it warms the ground; sunlight also transfers energy from the Sun to the Earth).

P4.1 Energy Transfer

Moving objects and waves transfer energy from one location to another. They also transfer energy to objects during interactions (e.g., sunlight transfers energy to the ground when it warms the ground; sunlight also transfers energy from the Sun to the Earth).

P4.1 Energy Transfer

Moving objects and waves transfer energy from one location to another. They also transfer energy to objects during interactions (e.g., sunlight transfers energy to the ground when it warms the ground; sunlight also transfers energy from the Sun to the Earth).



P4.1A Account for and represent energy into and out of systems using energy transfer diagrams.



P4.1A Account for and represent energy into and out of systems using energy transfer diagrams.



P4.1A Account for and represent energy into and out of systems using energy transfer diagrams.



P4.1A Account for and represent energy into and out of systems using energy transfer diagrams.



P4.1A Account for and represent energy into and out of systems using energy transfer diagrams.



P4.1A Account for and represent energy into and out of systems using energy transfer diagrams.

P4.1 Energy Transfer

Moving objects and waves transfer energy from one location to another. They also transfer energy to objects during interactions (e.g., sunlight transfers energy to the ground when it warms the ground; sunlight also transfers energy from the Sun to the Earth).

P4.1 Energy Transfer

Moving objects and waves transfer energy from one location to another. They also transfer energy to objects during interactions (e.g., sunlight transfers energy to the ground when it warms the ground; sunlight also transfers energy from the Sun to the Earth).

P4.1 Energy Transfer

Moving objects and waves transfer energy from one location to another. They also transfer energy to objects during interactions (e.g., sunlight transfers energy to the ground when it warms the ground; sunlight also transfers energy from the Sun to the Earth).

P4.1 Energy Transfer

Moving objects and waves transfer energy from one location to another. They also transfer energy to objects during interactions (e.g., sunlight transfers energy to the ground when it warms the ground; sunlight also transfers energy from the Sun to the Earth).

P4.1 Energy Transfer

Moving objects and waves transfer energy from one location to another. They also transfer energy to objects during interactions (e.g., sunlight transfers energy to the ground when it warms the ground; sunlight also transfers energy from the Sun to the Earth).

P4.1 Energy Transfer

Moving objects and waves transfer energy from one location to another. They also transfer energy to objects during interactions (e.g., sunlight transfers energy to the ground when it warms the ground; sunlight also transfers energy from the Sun to the Earth).

E

P4.1B Explain instances of energy transfer by waves and objects in everyday activities (e.g., why the ground gets warm during the day, how you hear a distant sound, why it hurts when you are hit by a baseball).

E

P4.1B Explain instances of energy transfer by waves and objects in everyday activities (e.g., why the ground gets warm during the day, how you hear a distant sound, why it hurts when you are hit by a baseball).

E

P4.1B Explain instances of energy transfer by waves and objects in everyday activities (e.g., why the ground gets warm during the day, how you hear a distant sound, why it hurts when you are hit by a baseball).

E

P4.1B Explain instances of energy transfer by waves and objects in everyday activities (e.g., why the ground gets warm during the day, how you hear a distant sound, why it hurts when you are hit by a baseball).

E

P4.1B Explain instances of energy transfer by waves and objects in everyday activities (e.g., why the ground gets warm during the day, how you hear a distant sound, why it hurts when you are hit by a baseball).

E

P4.1B Explain instances of energy transfer by waves and objects in everyday activities (e.g., why the ground gets warm during the day, how you hear a distant sound, why it hurts when you are hit by a baseball).

P4.1 Energy Transfer

Moving objects and waves transfer energy from one location to another. They also transfer energy to objects during interactions (e.g., sunlight transfers energy to the ground when it warms the ground; sunlight also transfers energy from the Sun to the Earth).

P4.1 Energy Transfer

Moving objects and waves transfer energy from one location to another. They also transfer energy to objects during interactions (e.g., sunlight transfers energy to the ground when it warms the ground; sunlight also transfers energy from the Sun to the Earth).

P4.1 Energy Transfer

Moving objects and waves transfer energy from one location to another. They also transfer energy to objects during interactions (e.g., sunlight transfers energy to the ground when it warms the ground; sunlight also transfers energy from the Sun to the Earth).

P4.1 Energy Transfer

Moving objects and waves transfer energy from one location to another. They also transfer energy to objects during interactions (e.g., sunlight transfers energy to the ground when it warms the ground; sunlight also transfers energy from the Sun to the Earth).

P4.1 Energy Transfer

Moving objects and waves transfer energy from one location to another. They also transfer energy to objects during interactions (e.g., sunlight transfers energy to the ground when it warms the ground; sunlight also transfers energy from the Sun to the Earth).

P4.1 Energy Transfer

Moving objects and waves transfer energy from one location to another. They also transfer energy to objects during interactions (e.g., sunlight transfers energy to the ground when it warms the ground; sunlight also transfers energy from the Sun to the Earth).



P4.1c Explain why work has a more precise scientific meaning than the meaning of work in everyday language.



P4.1c Explain why work has a more precise scientific meaning than the meaning of work in everyday language.



P4.1c Explain why work has a more precise scientific meaning than the meaning of work in everyday language.



P4.1c Explain why work has a more precise scientific meaning than the meaning of work in everyday language.



P4.1c Explain why work has a more precise scientific meaning than the meaning of work in everyday language.



P4.1c Explain why work has a more precise scientific meaning than the meaning of work in everyday language.

P4.1x Energy Transfer — Work

Work is the amount of energy transferred during an interaction. In mechanical systems, work is the amount of energy transferred as an object is moved through a distance, $W = F d$, where d is in the same direction as F . The total work done on an object depends on the net force acting on the object and the object's displacement.

P4.1x Energy Transfer — Work

Work is the amount of energy transferred during an interaction. In mechanical systems, work is the amount of energy transferred as an object is moved through a distance, $W = F d$, where d is in the same direction as F . The total work done on an object depends on the net force acting on the object and the object's displacement.

P4.1x Energy Transfer — Work

Work is the amount of energy transferred during an interaction. In mechanical systems, work is the amount of energy transferred as an object is moved through a distance, $W = F d$, where d is in the same direction as F . The total work done on an object depends on the net force acting on the object and the object's displacement.

P4.1x Energy Transfer — Work

Work is the amount of energy transferred during an interaction. In mechanical systems, work is the amount of energy transferred as an object is moved through a distance, $W = F d$, where d is in the same direction as F . The total work done on an object depends on the net force acting on the object and the object's displacement.

P4.1x Energy Transfer — Work

Work is the amount of energy transferred during an interaction. In mechanical systems, work is the amount of energy transferred as an object is moved through a distance, $W = F d$, where d is in the same direction as F . The total work done on an object depends on the net force acting on the object and the object's displacement.

P4.1x Energy Transfer — Work

Work is the amount of energy transferred during an interaction. In mechanical systems, work is the amount of energy transferred as an object is moved through a distance, $W = F d$, where d is in the same direction as F . The total work done on an object depends on the net force acting on the object and the object's displacement.



P4.Id Calculate the amount of work done on an object that is moved from one position to another.



P4.Id Calculate the amount of work done on an object that is moved from one position to another.



P4.Id Calculate the amount of work done on an object that is moved from one position to another.



P4.Id Calculate the amount of work done on an object that is moved from one position to another.



P4.Id Calculate the amount of work done on an object that is moved from one position to another.



P4.Id Calculate the amount of work done on an object that is moved from one position to another.

P4.1x Energy Transfer — Work

Work is the amount of energy transferred during an interaction. In mechanical systems, work is the amount of energy transferred as an object is moved through a distance, $W = F d$, where d is in the same direction as F . The total work done on an object depends on the net force acting on the object and the object's displacement.

P4.1x Energy Transfer — Work

Work is the amount of energy transferred during an interaction. In mechanical systems, work is the amount of energy transferred as an object is moved through a distance, $W = F d$, where d is in the same direction as F . The total work done on an object depends on the net force acting on the object and the object's displacement.

P4.1x Energy Transfer — Work

Work is the amount of energy transferred during an interaction. In mechanical systems, work is the amount of energy transferred as an object is moved through a distance, $W = F d$, where d is in the same direction as F . The total work done on an object depends on the net force acting on the object and the object's displacement.

P4.1x Energy Transfer — Work

Work is the amount of energy transferred during an interaction. In mechanical systems, work is the amount of energy transferred as an object is moved through a distance, $W = F d$, where d is in the same direction as F . The total work done on an object depends on the net force acting on the object and the object's displacement.

P4.1x Energy Transfer — Work

Work is the amount of energy transferred during an interaction. In mechanical systems, work is the amount of energy transferred as an object is moved through a distance, $W = F d$, where d is in the same direction as F . The total work done on an object depends on the net force acting on the object and the object's displacement.

P4.1x Energy Transfer — Work

Work is the amount of energy transferred during an interaction. In mechanical systems, work is the amount of energy transferred as an object is moved through a distance, $W = F d$, where d is in the same direction as F . The total work done on an object depends on the net force acting on the object and the object's displacement.



P4.1e Using the formula for work, derive a formula for change in potential energy of an object lifted a distance h .



P4.1e Using the formula for work, derive a formula for change in potential energy of an object lifted a distance h .



P4.1e Using the formula for work, derive a formula for change in potential energy of an object lifted a distance h .



P4.1e Using the formula for work, derive a formula for change in potential energy of an object lifted a distance h .



P4.1e Using the formula for work, derive a formula for change in potential energy of an object lifted a distance h .



P4.1e Using the formula for work, derive a formula for change in potential energy of an object lifted a distance h .

P4.2 Energy Transformation

Energy is often transformed from one form to another. The amount of energy before a transformation is equal to the amount of energy after the transformation. In most energy transformations, some energy is converted to thermal energy.

P4.2 Energy Transformation

Energy is often transformed from one form to another. The amount of energy before a transformation is equal to the amount of energy after the transformation. In most energy transformations, some energy is converted to thermal energy.

P4.2 Energy Transformation

Energy is often transformed from one form to another. The amount of energy before a transformation is equal to the amount of energy after the transformation. In most energy transformations, some energy is converted to thermal energy.

P4.2 Energy Transformation

Energy is often transformed from one form to another. The amount of energy before a transformation is equal to the amount of energy after the transformation. In most energy transformations, some energy is converted to thermal energy.

P4.2 Energy Transformation

Energy is often transformed from one form to another. The amount of energy before a transformation is equal to the amount of energy after the transformation. In most energy transformations, some energy is converted to thermal energy.

P4.2 Energy Transformation

Energy is often transformed from one form to another. The amount of energy before a transformation is equal to the amount of energy after the transformation. In most energy transformations, some energy is converted to thermal energy.



P4.2A Account for and represent energy transfer and transformation in complex processes (interactions).



P4.2A Account for and represent energy transfer and transformation in complex processes (interactions).



P4.2A Account for and represent energy transfer and transformation in complex processes (interactions).



P4.2A Account for and represent energy transfer and transformation in complex processes (interactions).



P4.2A Account for and represent energy transfer and transformation in complex processes (interactions).



P4.2A Account for and represent energy transfer and transformation in complex processes (interactions).

P4.2 Energy Transformation

Energy is often transformed from one form to another. The amount of energy before a transformation is equal to the amount of energy after the transformation. In most energy transformations, some energy is converted to thermal energy.

P4.2 Energy Transformation

Energy is often transformed from one form to another. The amount of energy before a transformation is equal to the amount of energy after the transformation. In most energy transformations, some energy is converted to thermal energy.

P4.2 Energy Transformation

Energy is often transformed from one form to another. The amount of energy before a transformation is equal to the amount of energy after the transformation. In most energy transformations, some energy is converted to thermal energy.

P4.2 Energy Transformation

Energy is often transformed from one form to another. The amount of energy before a transformation is equal to the amount of energy after the transformation. In most energy transformations, some energy is converted to thermal energy.

P4.2 Energy Transformation

Energy is often transformed from one form to another. The amount of energy before a transformation is equal to the amount of energy after the transformation. In most energy transformations, some energy is converted to thermal energy.

P4.2 Energy Transformation

Energy is often transformed from one form to another. The amount of energy before a transformation is equal to the amount of energy after the transformation. In most energy transformations, some energy is converted to thermal energy.



P4.2B Name devices that transform specific types of energy into other types (e.g., a device that transforms electricity into motion).



P4.2B Name devices that transform specific types of energy into other types (e.g., a device that transforms electricity into motion).



P4.2B Name devices that transform specific types of energy into other types (e.g., a device that transforms electricity into motion).



P4.2B Name devices that transform specific types of energy into other types (e.g., a device that transforms electricity into motion).



P4.2B Name devices that transform specific types of energy into other types (e.g., a device that transforms electricity into motion).



P4.2B Name devices that transform specific types of energy into other types (e.g., a device that transforms electricity into motion).

P4.2 Energy Transformation

Energy is often transformed from one form to another. The amount of energy before a transformation is equal to the amount of energy after the transformation. In most energy transformations, some energy is converted to thermal energy.

P4.2 Energy Transformation

Energy is often transformed from one form to another. The amount of energy before a transformation is equal to the amount of energy after the transformation. In most energy transformations, some energy is converted to thermal energy.

P4.2 Energy Transformation

Energy is often transformed from one form to another. The amount of energy before a transformation is equal to the amount of energy after the transformation. In most energy transformations, some energy is converted to thermal energy.

P4.2 Energy Transformation

Energy is often transformed from one form to another. The amount of energy before a transformation is equal to the amount of energy after the transformation. In most energy transformations, some energy is converted to thermal energy.

P4.2 Energy Transformation

Energy is often transformed from one form to another. The amount of energy before a transformation is equal to the amount of energy after the transformation. In most energy transformations, some energy is converted to thermal energy.

P4.2 Energy Transformation

Energy is often transformed from one form to another. The amount of energy before a transformation is equal to the amount of energy after the transformation. In most energy transformations, some energy is converted to thermal energy.



P4.2C Explain how energy is conserved in common systems (e.g., light incident on a transparent material, light incident on a leaf, mechanical energy in a collision).



P4.2C Explain how energy is conserved in common systems (e.g., light incident on a transparent material, light incident on a leaf, mechanical energy in a collision).



P4.2C Explain how energy is conserved in common systems (e.g., light incident on a transparent material, light incident on a leaf, mechanical energy in a collision).



P4.2C Explain how energy is conserved in common systems (e.g., light incident on a transparent material, light incident on a leaf, mechanical energy in a collision).



P4.2C Explain how energy is conserved in common systems (e.g., light incident on a transparent material, light incident on a leaf, mechanical energy in a collision).



P4.2C Explain how energy is conserved in common systems (e.g., light incident on a transparent material, light incident on a leaf, mechanical energy in a collision).

P4.2 Energy Transformation

Energy is often transformed from one form to another. The amount of energy before a transformation is equal to the amount of energy after the transformation. In most energy transformations, some energy is converted to thermal energy.

P4.2 Energy Transformation

Energy is often transformed from one form to another. The amount of energy before a transformation is equal to the amount of energy after the transformation. In most energy transformations, some energy is converted to thermal energy.

P4.2 Energy Transformation

Energy is often transformed from one form to another. The amount of energy before a transformation is equal to the amount of energy after the transformation. In most energy transformations, some energy is converted to thermal energy.

P4.2 Energy Transformation

Energy is often transformed from one form to another. The amount of energy before a transformation is equal to the amount of energy after the transformation. In most energy transformations, some energy is converted to thermal energy.

P4.2 Energy Transformation

Energy is often transformed from one form to another. The amount of energy before a transformation is equal to the amount of energy after the transformation. In most energy transformations, some energy is converted to thermal energy.

P4.2 Energy Transformation

Energy is often transformed from one form to another. The amount of energy before a transformation is equal to the amount of energy after the transformation. In most energy transformations, some energy is converted to thermal energy.

E

P4.2D Explain why all the stored energy in gasoline does not transform to mechanical energy of a vehicle.

E

P4.2D Explain why all the stored energy in gasoline does not transform to mechanical energy of a vehicle.

E

P4.2D Explain why all the stored energy in gasoline does not transform to mechanical energy of a vehicle.

E

P4.2D Explain why all the stored energy in gasoline does not transform to mechanical energy of a vehicle.

E

P4.2D Explain why all the stored energy in gasoline does not transform to mechanical energy of a vehicle.

E

P4.2D Explain why all the stored energy in gasoline does not transform to mechanical energy of a vehicle.

P4.2 Energy Transformation

Energy is often transformed from one form to another. The amount of energy before a transformation is equal to the amount of energy after the transformation. In most energy transformations, some energy is converted to thermal energy.

P4.2 Energy Transformation

Energy is often transformed from one form to another. The amount of energy before a transformation is equal to the amount of energy after the transformation. In most energy transformations, some energy is converted to thermal energy.

P4.2 Energy Transformation

Energy is often transformed from one form to another. The amount of energy before a transformation is equal to the amount of energy after the transformation. In most energy transformations, some energy is converted to thermal energy.

P4.2 Energy Transformation

Energy is often transformed from one form to another. The amount of energy before a transformation is equal to the amount of energy after the transformation. In most energy transformations, some energy is converted to thermal energy.

P4.2 Energy Transformation

Energy is often transformed from one form to another. The amount of energy before a transformation is equal to the amount of energy after the transformation. In most energy transformations, some energy is converted to thermal energy.

P4.2 Energy Transformation

Energy is often transformed from one form to another. The amount of energy before a transformation is equal to the amount of energy after the transformation. In most energy transformations, some energy is converted to thermal energy.



P4.2e Explain the energy transformation as an object (e.g., skydiver) falls at a steady velocity.



P4.2e Explain the energy transformation as an object (e.g., skydiver) falls at a steady velocity.



P4.2e Explain the energy transformation as an object (e.g., skydiver) falls at a steady velocity.



P4.2e Explain the energy transformation as an object (e.g., skydiver) falls at a steady velocity.



P4.2e Explain the energy transformation as an object (e.g., skydiver) falls at a steady velocity.



P4.2e Explain the energy transformation as an object (e.g., skydiver) falls at a steady velocity.

P4.2 Energy Transformation

Energy is often transformed from one form to another. The amount of energy before a transformation is equal to the amount of energy after the transformation. In most energy transformations, some energy is converted to thermal energy.

P4.2 Energy Transformation

Energy is often transformed from one form to another. The amount of energy before a transformation is equal to the amount of energy after the transformation. In most energy transformations, some energy is converted to thermal energy.

P4.2 Energy Transformation

Energy is often transformed from one form to another. The amount of energy before a transformation is equal to the amount of energy after the transformation. In most energy transformations, some energy is converted to thermal energy.

P4.2 Energy Transformation

Energy is often transformed from one form to another. The amount of energy before a transformation is equal to the amount of energy after the transformation. In most energy transformations, some energy is converted to thermal energy.

P4.2 Energy Transformation

Energy is often transformed from one form to another. The amount of energy before a transformation is equal to the amount of energy after the transformation. In most energy transformations, some energy is converted to thermal energy.

P4.2 Energy Transformation

Energy is often transformed from one form to another. The amount of energy before a transformation is equal to the amount of energy after the transformation. In most energy transformations, some energy is converted to thermal energy.



P4.2f Identify and label the energy inputs, transformations, and outputs using qualitative or quantitative representations in simple technological systems (e.g., toaster, motor, hair dryer) to show energy conservation.



P4.2f Identify and label the energy inputs, transformations, and outputs using qualitative or quantitative representations in simple technological systems (e.g., toaster, motor, hair dryer) to show energy conservation.



P4.2f Identify and label the energy inputs, transformations, and outputs using qualitative or quantitative representations in simple technological systems (e.g., toaster, motor, hair dryer) to show energy conservation.



P4.2f Identify and label the energy inputs, transformations, and outputs using qualitative or quantitative representations in simple technological systems (e.g., toaster, motor, hair dryer) to show energy conservation.



P4.2f Identify and label the energy inputs, transformations, and outputs using qualitative or quantitative representations in simple technological systems (e.g., toaster, motor, hair dryer) to show energy conservation.



P4.2f Identify and label the energy inputs, transformations, and outputs using qualitative or quantitative representations in simple technological systems (e.g., toaster, motor, hair dryer) to show energy conservation.

P4.3 Kinetic and Potential Energy

Moving objects have kinetic energy. Objects experiencing a force may have potential energy due to their relative positions (e.g., lifting an object or stretching a spring, energy stored in chemical bonds). Conversions between kinetic and gravitational potential energy are common in moving objects. In frictionless systems, the decrease in gravitational potential energy is equal to the increase in kinetic energy or vice versa.

P4.3 Kinetic and Potential Energy

Moving objects have kinetic energy. Objects experiencing a force may have potential energy due to their relative positions (e.g., lifting an object or stretching a spring, energy stored in chemical bonds). Conversions between kinetic and gravitational potential energy are common in moving objects. In frictionless systems, the decrease in gravitational potential energy is equal to the increase in kinetic energy or vice versa.

P4.3 Kinetic and Potential Energy

Moving objects have kinetic energy. Objects experiencing a force may have potential energy due to their relative positions (e.g., lifting an object or stretching a spring, energy stored in chemical bonds). Conversions between kinetic and gravitational potential energy are common in moving objects. In frictionless systems, the decrease in gravitational potential energy is equal to the increase in kinetic energy or vice versa.

P4.3 Kinetic and Potential Energy

Moving objects have kinetic energy. Objects experiencing a force may have potential energy due to their relative positions (e.g., lifting an object or stretching a spring, energy stored in chemical bonds). Conversions between kinetic and gravitational potential energy are common in moving objects. In frictionless systems, the decrease in gravitational potential energy is equal to the increase in kinetic energy or vice versa.

P4.3 Kinetic and Potential Energy

Moving objects have kinetic energy. Objects experiencing a force may have potential energy due to their relative positions (e.g., lifting an object or stretching a spring, energy stored in chemical bonds). Conversions between kinetic and gravitational potential energy are common in moving objects. In frictionless systems, the decrease in gravitational potential energy is equal to the increase in kinetic energy or vice versa.

P4.3 Kinetic and Potential Energy

Moving objects have kinetic energy. Objects experiencing a force may have potential energy due to their relative positions (e.g., lifting an object or stretching a spring, energy stored in chemical bonds). Conversions between kinetic and gravitational potential energy are common in moving objects. In frictionless systems, the decrease in gravitational potential energy is equal to the increase in kinetic energy or vice versa.



P4.3A Identify the form of energy in given situations (e.g., moving objects, stretched springs, rocks on cliffs, energy in food).



P4.3A Identify the form of energy in given situations (e.g., moving objects, stretched springs, rocks on cliffs, energy in food).



P4.3A Identify the form of energy in given situations (e.g., moving objects, stretched springs, rocks on cliffs, energy in food).



P4.3A Identify the form of energy in given situations (e.g., moving objects, stretched springs, rocks on cliffs, energy in food).



P4.3A Identify the form of energy in given situations (e.g., moving objects, stretched springs, rocks on cliffs, energy in food).



P4.3A Identify the form of energy in given situations (e.g., moving objects, stretched springs, rocks on cliffs, energy in food).

P4.3 Kinetic and Potential Energy

Moving objects have kinetic energy. Objects experiencing a force may have potential energy due to their relative positions (e.g., lifting an object or stretching a spring, energy stored in chemical bonds). Conversions between kinetic and gravitational potential energy are common in moving objects. In frictionless systems, the decrease in gravitational potential energy is equal to the increase in kinetic energy or vice versa.

P4.3 Kinetic and Potential Energy

Moving objects have kinetic energy. Objects experiencing a force may have potential energy due to their relative positions (e.g., lifting an object or stretching a spring, energy stored in chemical bonds). Conversions between kinetic and gravitational potential energy are common in moving objects. In frictionless systems, the decrease in gravitational potential energy is equal to the increase in kinetic energy or vice versa.

P4.3 Kinetic and Potential Energy

Moving objects have kinetic energy. Objects experiencing a force may have potential energy due to their relative positions (e.g., lifting an object or stretching a spring, energy stored in chemical bonds). Conversions between kinetic and gravitational potential energy are common in moving objects. In frictionless systems, the decrease in gravitational potential energy is equal to the increase in kinetic energy or vice versa.

P4.3 Kinetic and Potential Energy

Moving objects have kinetic energy. Objects experiencing a force may have potential energy due to their relative positions (e.g., lifting an object or stretching a spring, energy stored in chemical bonds). Conversions between kinetic and gravitational potential energy are common in moving objects. In frictionless systems, the decrease in gravitational potential energy is equal to the increase in kinetic energy or vice versa.

P4.3 Kinetic and Potential Energy

Moving objects have kinetic energy. Objects experiencing a force may have potential energy due to their relative positions (e.g., lifting an object or stretching a spring, energy stored in chemical bonds). Conversions between kinetic and gravitational potential energy are common in moving objects. In frictionless systems, the decrease in gravitational potential energy is equal to the increase in kinetic energy or vice versa.

P4.3 Kinetic and Potential Energy

Moving objects have kinetic energy. Objects experiencing a force may have potential energy due to their relative positions (e.g., lifting an object or stretching a spring, energy stored in chemical bonds). Conversions between kinetic and gravitational potential energy are common in moving objects. In frictionless systems, the decrease in gravitational potential energy is equal to the increase in kinetic energy or vice versa.



P4.3B Describe the transformation between potential and kinetic energy in simple mechanical systems (e.g., pendulums, roller coasters, ski lifts).



P4.3B Describe the transformation between potential and kinetic energy in simple mechanical systems (e.g., pendulums, roller coasters, ski lifts).



P4.3B Describe the transformation between potential and kinetic energy in simple mechanical systems (e.g., pendulums, roller coasters, ski lifts).



P4.3B Describe the transformation between potential and kinetic energy in simple mechanical systems (e.g., pendulums, roller coasters, ski lifts).



P4.3B Describe the transformation between potential and kinetic energy in simple mechanical systems (e.g., pendulums, roller coasters, ski lifts).



P4.3B Describe the transformation between potential and kinetic energy in simple mechanical systems (e.g., pendulums, roller coasters, ski lifts).

P4.3 Kinetic and Potential Energy

Moving objects have kinetic energy. Objects experiencing a force may have potential energy due to their relative positions (e.g., lifting an object or stretching a spring, energy stored in chemical bonds). Conversions between kinetic and gravitational potential energy are common in moving objects. In frictionless systems, the decrease in gravitational potential energy is equal to the increase in kinetic energy or vice versa.

P4.3 Kinetic and Potential Energy

Moving objects have kinetic energy. Objects experiencing a force may have potential energy due to their relative positions (e.g., lifting an object or stretching a spring, energy stored in chemical bonds). Conversions between kinetic and gravitational potential energy are common in moving objects. In frictionless systems, the decrease in gravitational potential energy is equal to the increase in kinetic energy or vice versa.

P4.3 Kinetic and Potential Energy

Moving objects have kinetic energy. Objects experiencing a force may have potential energy due to their relative positions (e.g., lifting an object or stretching a spring, energy stored in chemical bonds). Conversions between kinetic and gravitational potential energy are common in moving objects. In frictionless systems, the decrease in gravitational potential energy is equal to the increase in kinetic energy or vice versa.

P4.3 Kinetic and Potential Energy

Moving objects have kinetic energy. Objects experiencing a force may have potential energy due to their relative positions (e.g., lifting an object or stretching a spring, energy stored in chemical bonds). Conversions between kinetic and gravitational potential energy are common in moving objects. In frictionless systems, the decrease in gravitational potential energy is equal to the increase in kinetic energy or vice versa.

P4.3 Kinetic and Potential Energy

Moving objects have kinetic energy. Objects experiencing a force may have potential energy due to their relative positions (e.g., lifting an object or stretching a spring, energy stored in chemical bonds). Conversions between kinetic and gravitational potential energy are common in moving objects. In frictionless systems, the decrease in gravitational potential energy is equal to the increase in kinetic energy or vice versa.

P4.3 Kinetic and Potential Energy

Moving objects have kinetic energy. Objects experiencing a force may have potential energy due to their relative positions (e.g., lifting an object or stretching a spring, energy stored in chemical bonds). Conversions between kinetic and gravitational potential energy are common in moving objects. In frictionless systems, the decrease in gravitational potential energy is equal to the increase in kinetic energy or vice versa.

E

P4.3C Explain why all mechanical systems require an external energy source to maintain their motion

E

P4.3C Explain why all mechanical systems require an external energy source to maintain their motion

E

P4.3C Explain why all mechanical systems require an external energy source to maintain their motion

E

P4.3C Explain why all mechanical systems require an external energy source to maintain their motion

E

P4.3C Explain why all mechanical systems require an external energy source to maintain their motion

E

P4.3C Explain why all mechanical systems require an external energy source to maintain their motion

P4.3x Kinetic and Potential Energy — Calculations

The kinetic energy of an object is related to the mass of an object and its speed: $KE = 1/2 mv^2$.

P4.3x Kinetic and Potential Energy — Calculations

The kinetic energy of an object is related to the mass of an object and its speed: $KE = 1/2 mv^2$.

P4.3x Kinetic and Potential Energy — Calculations

The kinetic energy of an object is related to the mass of an object and its speed: $KE = 1/2 mv^2$.

P4.3x Kinetic and Potential Energy — Calculations

The kinetic energy of an object is related to the mass of an object and its speed: $KE = 1/2 mv^2$.

P4.3x Kinetic and Potential Energy — Calculations

The kinetic energy of an object is related to the mass of an object and its speed: $KE = 1/2 mv^2$.

P4.3x Kinetic and Potential Energy — Calculations

The kinetic energy of an object is related to the mass of an object and its speed: $KE = 1/2 mv^2$.



P4.3d Rank the amount of kinetic energy from highest to lowest of everyday examples of moving objects.



P4.3d Rank the amount of kinetic energy from highest to lowest of everyday examples of moving objects.



P4.3d Rank the amount of kinetic energy from highest to lowest of everyday examples of moving objects.



P4.3d Rank the amount of kinetic energy from highest to lowest of everyday examples of moving objects.



P4.3d Rank the amount of kinetic energy from highest to lowest of everyday examples of moving objects.



P4.3d Rank the amount of kinetic energy from highest to lowest of everyday examples of moving objects.

P4.3x Kinetic and Potential Energy — Calculations

The kinetic energy of an object is related to the mass of an object and its speed: $KE = 1/2 mv^2$.

P4.3x Kinetic and Potential Energy — Calculations

The kinetic energy of an object is related to the mass of an object and its speed: $KE = 1/2 mv^2$.

P4.3x Kinetic and Potential Energy — Calculations

The kinetic energy of an object is related to the mass of an object and its speed: $KE = 1/2 mv^2$.

P4.3x Kinetic and Potential Energy — Calculations

The kinetic energy of an object is related to the mass of an object and its speed: $KE = 1/2 mv^2$.

P4.3x Kinetic and Potential Energy — Calculations

The kinetic energy of an object is related to the mass of an object and its speed: $KE = 1/2 mv^2$.

P4.3x Kinetic and Potential Energy — Calculations

The kinetic energy of an object is related to the mass of an object and its speed: $KE = 1/2 mv^2$.

E

P4.3e Calculate the changes in kinetic and potential energy in simple mechanical systems (e.g., pendulums, roller coasters, ski lifts) using the formulas for kinetic energy and potential energy.

E

P4.3e Calculate the changes in kinetic and potential energy in simple mechanical systems (e.g., pendulums, roller coasters, ski lifts) using the formulas for kinetic energy and potential energy.

E

P4.3e Calculate the changes in kinetic and potential energy in simple mechanical systems (e.g., pendulums, roller coasters, ski lifts) using the formulas for kinetic energy and potential energy.

E

P4.3e Calculate the changes in kinetic and potential energy in simple mechanical systems (e.g., pendulums, roller coasters, ski lifts) using the formulas for kinetic energy and potential energy.

E

P4.3e Calculate the changes in kinetic and potential energy in simple mechanical systems (e.g., pendulums, roller coasters, ski lifts) using the formulas for kinetic energy and potential energy.

E

P4.3e Calculate the changes in kinetic and potential energy in simple mechanical systems (e.g., pendulums, roller coasters, ski lifts) using the formulas for kinetic energy and potential energy.

P4.3x Kinetic and Potential Energy — Calculations

The kinetic energy of an object is related to the mass of an object and its speed: $KE = 1/2 mv^2$.

P4.3x Kinetic and Potential Energy — Calculations

The kinetic energy of an object is related to the mass of an object and its speed: $KE = 1/2 mv^2$.

P4.3x Kinetic and Potential Energy — Calculations

The kinetic energy of an object is related to the mass of an object and its speed: $KE = 1/2 mv^2$.

P4.3x Kinetic and Potential Energy — Calculations

The kinetic energy of an object is related to the mass of an object and its speed: $KE = 1/2 mv^2$.

P4.3x Kinetic and Potential Energy — Calculations

The kinetic energy of an object is related to the mass of an object and its speed: $KE = 1/2 mv^2$.

P4.3x Kinetic and Potential Energy — Calculations

The kinetic energy of an object is related to the mass of an object and its speed: $KE = 1/2 mv^2$.



P4.3f Calculate the impact speed (ignoring air resistance) of an object dropped from a specific height or the maximum height reached by an object (ignoring air resistance), given the initial vertical velocity.



P4.3f Calculate the impact speed (ignoring air resistance) of an object dropped from a specific height or the maximum height reached by an object (ignoring air resistance), given the initial vertical velocity.



P4.3f Calculate the impact speed (ignoring air resistance) of an object dropped from a specific height or the maximum height reached by an object (ignoring air resistance), given the initial vertical velocity.



P4.3f Calculate the impact speed (ignoring air resistance) of an object dropped from a specific height or the maximum height reached by an object (ignoring air resistance), given the initial vertical velocity.



P4.3f Calculate the impact speed (ignoring air resistance) of an object dropped from a specific height or the maximum height reached by an object (ignoring air resistance), given the initial vertical velocity.



P4.3f Calculate the impact speed (ignoring air resistance) of an object dropped from a specific height or the maximum height reached by an object (ignoring air resistance), given the initial vertical velocity.

P4.4 Wave Characteristics

Waves (mechanical and electromagnetic) are described by their wavelength, amplitude, frequency, and speed.

P4.4 Wave Characteristics

Waves (mechanical and electromagnetic) are described by their wavelength, amplitude, frequency, and speed.

P4.4 Wave Characteristics

Waves (mechanical and electromagnetic) are described by their wavelength, amplitude, frequency, and speed.

P4.4 Wave Characteristics

Waves (mechanical and electromagnetic) are described by their wavelength, amplitude, frequency, and speed.

P4.4 Wave Characteristics

Waves (mechanical and electromagnetic) are described by their wavelength, amplitude, frequency, and speed.

P4.4 Wave Characteristics

Waves (mechanical and electromagnetic) are described by their wavelength, amplitude, frequency, and speed.

E

P4.4A Describe specific mechanical waves (e.g., on a demonstration spring, on the ocean) in terms of wavelength, amplitude, frequency, and speed.

E

P4.4A Describe specific mechanical waves (e.g., on a demonstration spring, on the ocean) in terms of wavelength, amplitude, frequency, and speed.

E

P4.4A Describe specific mechanical waves (e.g., on a demonstration spring, on the ocean) in terms of wavelength, amplitude, frequency, and speed.

E

P4.4A Describe specific mechanical waves (e.g., on a demonstration spring, on the ocean) in terms of wavelength, amplitude, frequency, and speed.

E

P4.4A Describe specific mechanical waves (e.g., on a demonstration spring, on the ocean) in terms of wavelength, amplitude, frequency, and speed.

E

P4.4A Describe specific mechanical waves (e.g., on a demonstration spring, on the ocean) in terms of wavelength, amplitude, frequency, and speed.

P4.4 Wave Characteristics

Waves (mechanical and electromagnetic) are described by their wavelength, amplitude, frequency, and speed.

P4.4 Wave Characteristics

Waves (mechanical and electromagnetic) are described by their wavelength, amplitude, frequency, and speed.

P4.4 Wave Characteristics

Waves (mechanical and electromagnetic) are described by their wavelength, amplitude, frequency, and speed.

P4.4 Wave Characteristics

Waves (mechanical and electromagnetic) are described by their wavelength, amplitude, frequency, and speed.

P4.4 Wave Characteristics

Waves (mechanical and electromagnetic) are described by their wavelength, amplitude, frequency, and speed.

P4.4 Wave Characteristics

Waves (mechanical and electromagnetic) are described by their wavelength, amplitude, frequency, and speed.



P4.4B Identify everyday examples of transverse and compression (longitudinal) waves.



P4.4B Identify everyday examples of transverse and compression (longitudinal) waves.



P4.4B Identify everyday examples of transverse and compression (longitudinal) waves.



P4.4B Identify everyday examples of transverse and compression (longitudinal) waves.



P4.4B Identify everyday examples of transverse and compression (longitudinal) waves.



P4.4B Identify everyday examples of transverse and compression (longitudinal) waves.

P4.4 Wave Characteristics

Waves (mechanical and electromagnetic) are described by their wavelength, amplitude, frequency, and speed.

P4.4 Wave Characteristics

Waves (mechanical and electromagnetic) are described by their wavelength, amplitude, frequency, and speed.

P4.4 Wave Characteristics

Waves (mechanical and electromagnetic) are described by their wavelength, amplitude, frequency, and speed.

P4.4 Wave Characteristics

Waves (mechanical and electromagnetic) are described by their wavelength, amplitude, frequency, and speed.

P4.4 Wave Characteristics

Waves (mechanical and electromagnetic) are described by their wavelength, amplitude, frequency, and speed.

P4.4 Wave Characteristics

Waves (mechanical and electromagnetic) are described by their wavelength, amplitude, frequency, and speed.



P4.4C Compare and contrast transverse and compression (longitudinal) waves in terms of wavelength, amplitude, and frequency.



P4.4C Compare and contrast transverse and compression (longitudinal) waves in terms of wavelength, amplitude, and frequency.



P4.4C Compare and contrast transverse and compression (longitudinal) waves in terms of wavelength, amplitude, and frequency.



P4.4C Compare and contrast transverse and compression (longitudinal) waves in terms of wavelength, amplitude, and frequency.



P4.4C Compare and contrast transverse and compression (longitudinal) waves in terms of wavelength, amplitude, and frequency.



P4.4C Compare and contrast transverse and compression (longitudinal) waves in terms of wavelength, amplitude, and frequency.

P4.4x Wave Characteristics — Calculations

Wave velocity, wavelength, and frequency are related by $v = \lambda f$. The energy transferred by a wave is proportional to the square of the amplitude of vibration and its frequency.

P4.4x Wave Characteristics — Calculations

Wave velocity, wavelength, and frequency are related by $v = \lambda f$. The energy transferred by a wave is proportional to the square of the amplitude of vibration and its frequency.

P4.4x Wave Characteristics — Calculations

Wave velocity, wavelength, and frequency are related by $v = \lambda f$. The energy transferred by a wave is proportional to the square of the amplitude of vibration and its frequency.

P4.4x Wave Characteristics — Calculations

Wave velocity, wavelength, and frequency are related by $v = \lambda f$. The energy transferred by a wave is proportional to the square of the amplitude of vibration and its frequency.

P4.4x Wave Characteristics — Calculations

Wave velocity, wavelength, and frequency are related by $v = \lambda f$. The energy transferred by a wave is proportional to the square of the amplitude of vibration and its frequency.

P4.4x Wave Characteristics — Calculations

Wave velocity, wavelength, and frequency are related by $v = \lambda f$. The energy transferred by a wave is proportional to the square of the amplitude of vibration and its frequency.



P4.4d Demonstrate that frequency and wavelength of a wave are inversely proportional in a given medium.



P4.4d Demonstrate that frequency and wavelength of a wave are inversely proportional in a given medium.



P4.4d Demonstrate that frequency and wavelength of a wave are inversely proportional in a given medium.



P4.4d Demonstrate that frequency and wavelength of a wave are inversely proportional in a given medium.



P4.4d Demonstrate that frequency and wavelength of a wave are inversely proportional in a given medium.



P4.4d Demonstrate that frequency and wavelength of a wave are inversely proportional in a given medium.

P4.4x Wave Characteristics — Calculations

Wave velocity, wavelength, and frequency are related by $v = \lambda f$. The energy transferred by a wave is proportional to the square of the amplitude of vibration and its frequency.

P4.4x Wave Characteristics — Calculations

Wave velocity, wavelength, and frequency are related by $v = \lambda f$. The energy transferred by a wave is proportional to the square of the amplitude of vibration and its frequency.

P4.4x Wave Characteristics — Calculations

Wave velocity, wavelength, and frequency are related by $v = \lambda f$. The energy transferred by a wave is proportional to the square of the amplitude of vibration and its frequency.

P4.4x Wave Characteristics — Calculations

Wave velocity, wavelength, and frequency are related by $v = \lambda f$. The energy transferred by a wave is proportional to the square of the amplitude of vibration and its frequency.

P4.4x Wave Characteristics — Calculations

Wave velocity, wavelength, and frequency are related by $v = \lambda f$. The energy transferred by a wave is proportional to the square of the amplitude of vibration and its frequency.

P4.4x Wave Characteristics — Calculations

Wave velocity, wavelength, and frequency are related by $v = \lambda f$. The energy transferred by a wave is proportional to the square of the amplitude of vibration and its frequency.



P4.4e Calculate the amount of energy transferred by transverse or compression waves of different amplitudes and frequencies (e.g., seismic waves).



P4.4e Calculate the amount of energy transferred by transverse or compression waves of different amplitudes and frequencies (e.g., seismic waves).



P4.4e Calculate the amount of energy transferred by transverse or compression waves of different amplitudes and frequencies (e.g., seismic waves).



P4.4e Calculate the amount of energy transferred by transverse or compression waves of different amplitudes and frequencies (e.g., seismic waves).



P4.4e Calculate the amount of energy transferred by transverse or compression waves of different amplitudes and frequencies (e.g., seismic waves).



P4.4e Calculate the amount of energy transferred by transverse or compression waves of different amplitudes and frequencies (e.g., seismic waves).

P4.5 Mechanical Wave Propagation

Vibrations in matter initiate mechanical waves (e.g., water waves, sound waves, seismic waves), which may propagate in all directions and decrease in intensity in proportion to the distance squared for a point source. Waves transfer energy from one place to another without transferring mass.

P4.5 Mechanical Wave Propagation

Vibrations in matter initiate mechanical waves (e.g., water waves, sound waves, seismic waves), which may propagate in all directions and decrease in intensity in proportion to the distance squared for a point source. Waves transfer energy from one place to another without transferring mass.

P4.5 Mechanical Wave Propagation

Vibrations in matter initiate mechanical waves (e.g., water waves, sound waves, seismic waves), which may propagate in all directions and decrease in intensity in proportion to the distance squared for a point source. Waves transfer energy from one place to another without transferring mass.

P4.5 Mechanical Wave Propagation

Vibrations in matter initiate mechanical waves (e.g., water waves, sound waves, seismic waves), which may propagate in all directions and decrease in intensity in proportion to the distance squared for a point source. Waves transfer energy from one place to another without transferring mass.

P4.5 Mechanical Wave Propagation

Vibrations in matter initiate mechanical waves (e.g., water waves, sound waves, seismic waves), which may propagate in all directions and decrease in intensity in proportion to the distance squared for a point source. Waves transfer energy from one place to another without transferring mass.

P4.5 Mechanical Wave Propagation

Vibrations in matter initiate mechanical waves (e.g., water waves, sound waves, seismic waves), which may propagate in all directions and decrease in intensity in proportion to the distance squared for a point source. Waves transfer energy from one place to another without transferring mass.



P4.5A Identify everyday examples of energy transfer by waves and their sources.



P4.5A Identify everyday examples of energy transfer by waves and their sources.



P4.5A Identify everyday examples of energy transfer by waves and their sources.



P4.5A Identify everyday examples of energy transfer by waves and their sources.



P4.5A Identify everyday examples of energy transfer by waves and their sources.



P4.5A Identify everyday examples of energy transfer by waves and their sources.

P4.5 Mechanical Wave Propagation

Vibrations in matter initiate mechanical waves (e.g., water waves, sound waves, seismic waves), which may propagate in all directions and decrease in intensity in proportion to the distance squared for a point source. Waves transfer energy from one place to another without transferring mass.

P4.5 Mechanical Wave Propagation

Vibrations in matter initiate mechanical waves (e.g., water waves, sound waves, seismic waves), which may propagate in all directions and decrease in intensity in proportion to the distance squared for a point source. Waves transfer energy from one place to another without transferring mass.

P4.5 Mechanical Wave Propagation

Vibrations in matter initiate mechanical waves (e.g., water waves, sound waves, seismic waves), which may propagate in all directions and decrease in intensity in proportion to the distance squared for a point source. Waves transfer energy from one place to another without transferring mass.

P4.5 Mechanical Wave Propagation

Vibrations in matter initiate mechanical waves (e.g., water waves, sound waves, seismic waves), which may propagate in all directions and decrease in intensity in proportion to the distance squared for a point source. Waves transfer energy from one place to another without transferring mass.

P4.5 Mechanical Wave Propagation

Vibrations in matter initiate mechanical waves (e.g., water waves, sound waves, seismic waves), which may propagate in all directions and decrease in intensity in proportion to the distance squared for a point source. Waves transfer energy from one place to another without transferring mass.

P4.5 Mechanical Wave Propagation

Vibrations in matter initiate mechanical waves (e.g., water waves, sound waves, seismic waves), which may propagate in all directions and decrease in intensity in proportion to the distance squared for a point source. Waves transfer energy from one place to another without transferring mass.

E

P4.5B Explain why an object (e.g., fishing bobber) does not move forward as a wave passes under it.

E

P4.5B Explain why an object (e.g., fishing bobber) does not move forward as a wave passes under it.

E

P4.5B Explain why an object (e.g., fishing bobber) does not move forward as a wave passes under it.

E

P4.5B Explain why an object (e.g., fishing bobber) does not move forward as a wave passes under it.

E

P4.5B Explain why an object (e.g., fishing bobber) does not move forward as a wave passes under it.

E

P4.5B Explain why an object (e.g., fishing bobber) does not move forward as a wave passes under it.

P4.5 Mechanical Wave Propagation

Vibrations in matter initiate mechanical waves (e.g., water waves, sound waves, seismic waves), which may propagate in all directions and decrease in intensity in proportion to the distance squared for a point source. Waves transfer energy from one place to another without transferring mass.

P4.5 Mechanical Wave Propagation

Vibrations in matter initiate mechanical waves (e.g., water waves, sound waves, seismic waves), which may propagate in all directions and decrease in intensity in proportion to the distance squared for a point source. Waves transfer energy from one place to another without transferring mass.

P4.5 Mechanical Wave Propagation

Vibrations in matter initiate mechanical waves (e.g., water waves, sound waves, seismic waves), which may propagate in all directions and decrease in intensity in proportion to the distance squared for a point source. Waves transfer energy from one place to another without transferring mass.

P4.5 Mechanical Wave Propagation

Vibrations in matter initiate mechanical waves (e.g., water waves, sound waves, seismic waves), which may propagate in all directions and decrease in intensity in proportion to the distance squared for a point source. Waves transfer energy from one place to another without transferring mass.

P4.5 Mechanical Wave Propagation

Vibrations in matter initiate mechanical waves (e.g., water waves, sound waves, seismic waves), which may propagate in all directions and decrease in intensity in proportion to the distance squared for a point source. Waves transfer energy from one place to another without transferring mass.

P4.5 Mechanical Wave Propagation

Vibrations in matter initiate mechanical waves (e.g., water waves, sound waves, seismic waves), which may propagate in all directions and decrease in intensity in proportion to the distance squared for a point source. Waves transfer energy from one place to another without transferring mass.

E

P4.5C Provide evidence to support the claim that sound is energy transferred by a wave, not energy transferred by particles.

E

P4.5C Provide evidence to support the claim that sound is energy transferred by a wave, not energy transferred by particles.

E

P4.5C Provide evidence to support the claim that sound is energy transferred by a wave, not energy transferred by particles.

E

P4.5C Provide evidence to support the claim that sound is energy transferred by a wave, not energy transferred by particles.

E

P4.5C Provide evidence to support the claim that sound is energy transferred by a wave, not energy transferred by particles.

E

P4.5C Provide evidence to support the claim that sound is energy transferred by a wave, not energy transferred by particles.

P4.5 Mechanical Wave Propagation

Vibrations in matter initiate mechanical waves (e.g., water waves, sound waves, seismic waves), which may propagate in all directions and decrease in intensity in proportion to the distance squared for a point source. Waves transfer energy from one place to another without transferring mass.

P4.5 Mechanical Wave Propagation

Vibrations in matter initiate mechanical waves (e.g., water waves, sound waves, seismic waves), which may propagate in all directions and decrease in intensity in proportion to the distance squared for a point source. Waves transfer energy from one place to another without transferring mass.

P4.5 Mechanical Wave Propagation

Vibrations in matter initiate mechanical waves (e.g., water waves, sound waves, seismic waves), which may propagate in all directions and decrease in intensity in proportion to the distance squared for a point source. Waves transfer energy from one place to another without transferring mass.

P4.5 Mechanical Wave Propagation

Vibrations in matter initiate mechanical waves (e.g., water waves, sound waves, seismic waves), which may propagate in all directions and decrease in intensity in proportion to the distance squared for a point source. Waves transfer energy from one place to another without transferring mass.

P4.5 Mechanical Wave Propagation

Vibrations in matter initiate mechanical waves (e.g., water waves, sound waves, seismic waves), which may propagate in all directions and decrease in intensity in proportion to the distance squared for a point source. Waves transfer energy from one place to another without transferring mass.

P4.5 Mechanical Wave Propagation

Vibrations in matter initiate mechanical waves (e.g., water waves, sound waves, seismic waves), which may propagate in all directions and decrease in intensity in proportion to the distance squared for a point source. Waves transfer energy from one place to another without transferring mass.

E

P4.5D Explain how waves propagate from vibrating sources and why the intensity decreases with the square of the distance from a point source.

E

P4.5D Explain how waves propagate from vibrating sources and why the intensity decreases with the square of the distance from a point source.

E

P4.5D Explain how waves propagate from vibrating sources and why the intensity decreases with the square of the distance from a point source.

E

P4.5D Explain how waves propagate from vibrating sources and why the intensity decreases with the square of the distance from a point source.

E

P4.5D Explain how waves propagate from vibrating sources and why the intensity decreases with the square of the distance from a point source.

E

P4.5D Explain how waves propagate from vibrating sources and why the intensity decreases with the square of the distance from a point source.

P4.5 Mechanical Wave Propagation

Vibrations in matter initiate mechanical waves (e.g., water waves, sound waves, seismic waves), which may propagate in all directions and decrease in intensity in proportion to the distance squared for a point source. Waves transfer energy from one place to another without transferring mass.

P4.5 Mechanical Wave Propagation

Vibrations in matter initiate mechanical waves (e.g., water waves, sound waves, seismic waves), which may propagate in all directions and decrease in intensity in proportion to the distance squared for a point source. Waves transfer energy from one place to another without transferring mass.

P4.5 Mechanical Wave Propagation

Vibrations in matter initiate mechanical waves (e.g., water waves, sound waves, seismic waves), which may propagate in all directions and decrease in intensity in proportion to the distance squared for a point source. Waves transfer energy from one place to another without transferring mass.

P4.5 Mechanical Wave Propagation

Vibrations in matter initiate mechanical waves (e.g., water waves, sound waves, seismic waves), which may propagate in all directions and decrease in intensity in proportion to the distance squared for a point source. Waves transfer energy from one place to another without transferring mass.

P4.5 Mechanical Wave Propagation

Vibrations in matter initiate mechanical waves (e.g., water waves, sound waves, seismic waves), which may propagate in all directions and decrease in intensity in proportion to the distance squared for a point source. Waves transfer energy from one place to another without transferring mass.

P4.5 Mechanical Wave Propagation

Vibrations in matter initiate mechanical waves (e.g., water waves, sound waves, seismic waves), which may propagate in all directions and decrease in intensity in proportion to the distance squared for a point source. Waves transfer energy from one place to another without transferring mass.

E

P4.5E Explain why everyone in a classroom can hear one person speaking, but why an amplification system is often used in the rear of a large concert auditorium.

E

P4.5E Explain why everyone in a classroom can hear one person speaking, but why an amplification system is often used in the rear of a large concert auditorium.

E

P4.5E Explain why everyone in a classroom can hear one person speaking, but why an amplification system is often used in the rear of a large concert auditorium.

E

P4.5E Explain why everyone in a classroom can hear one person speaking, but why an amplification system is often used in the rear of a large concert auditorium.

E

P4.5E Explain why everyone in a classroom can hear one person speaking, but why an amplification system is often used in the rear of a large concert auditorium.

E

P4.5E Explain why everyone in a classroom can hear one person speaking, but why an amplification system is often used in the rear of a large concert auditorium.

P4.6 Electromagnetic Waves

Electromagnetic waves (e.g., radio, microwave, infrared, visible light, ultraviolet, x-ray) are produced by changing the motion (acceleration) of charges or by changing magnetic fields. Electromagnetic waves can travel through matter, but they do not require a material medium. (That is, they also travel through empty space.) All electromagnetic waves move in a vacuum at the speed of light. Types of electromagnetic radiation are distinguished from each other by their wavelength and energy.

P4.6 Electromagnetic Waves

Electromagnetic waves (e.g., radio, microwave, infrared, visible light, ultraviolet, x-ray) are produced by changing the motion (acceleration) of charges or by changing magnetic fields. Electromagnetic waves can travel through matter, but they do not require a material medium. (That is, they also travel through empty space.) All electromagnetic waves move in a vacuum at the speed of light. Types of electromagnetic radiation are distinguished from each other by their wavelength and energy.

P4.6 Electromagnetic Waves

Electromagnetic waves (e.g., radio, microwave, infrared, visible light, ultraviolet, x-ray) are produced by changing the motion (acceleration) of charges or by changing magnetic fields. Electromagnetic waves can travel through matter, but they do not require a material medium. (That is, they also travel through empty space.) All electromagnetic waves move in a vacuum at the speed of light. Types of electromagnetic radiation are distinguished from each other by their wavelength and energy.

P4.6 Electromagnetic Waves

Electromagnetic waves (e.g., radio, microwave, infrared, visible light, ultraviolet, x-ray) are produced by changing the motion (acceleration) of charges or by changing magnetic fields. Electromagnetic waves can travel through matter, but they do not require a material medium. (That is, they also travel through empty space.) All electromagnetic waves move in a vacuum at the speed of light. Types of electromagnetic radiation are distinguished from each other by their wavelength and energy.

P4.6 Electromagnetic Waves

Electromagnetic waves (e.g., radio, microwave, infrared, visible light, ultraviolet, x-ray) are produced by changing the motion (acceleration) of charges or by changing magnetic fields. Electromagnetic waves can travel through matter, but they do not require a material medium. (That is, they also travel through empty space.) All electromagnetic waves move in a vacuum at the speed of light. Types of electromagnetic radiation are distinguished from each other by their wavelength and energy.

P4.6 Electromagnetic Waves

Electromagnetic waves (e.g., radio, microwave, infrared, visible light, ultraviolet, x-ray) are produced by changing the motion (acceleration) of charges or by changing magnetic fields. Electromagnetic waves can travel through matter, but they do not require a material medium. (That is, they also travel through empty space.) All electromagnetic waves move in a vacuum at the speed of light. Types of electromagnetic radiation are distinguished from each other by their wavelength and energy.



P4.6A Identify the different regions on the electromagnetic spectrum and compare them in terms of wavelength, frequency, and energy.



P4.6A Identify the different regions on the electromagnetic spectrum and compare them in terms of wavelength, frequency, and energy.



P4.6A Identify the different regions on the electromagnetic spectrum and compare them in terms of wavelength, frequency, and energy.



P4.6A Identify the different regions on the electromagnetic spectrum and compare them in terms of wavelength, frequency, and energy.



P4.6A Identify the different regions on the electromagnetic spectrum and compare them in terms of wavelength, frequency, and energy.



P4.6A Identify the different regions on the electromagnetic spectrum and compare them in terms of wavelength, frequency, and energy.

P4.6 Electromagnetic Waves

Electromagnetic waves (e.g., radio, microwave, infrared, visible light, ultraviolet, x-ray) are produced by changing the motion (acceleration) of charges or by changing magnetic fields. Electromagnetic waves can travel through matter, but they do not require a material medium. (That is, they also travel through empty space.) All electromagnetic waves move in a vacuum at the speed of light. Types of electromagnetic radiation are distinguished from each other by their wavelength and energy.

P4.6 Electromagnetic Waves

Electromagnetic waves (e.g., radio, microwave, infrared, visible light, ultraviolet, x-ray) are produced by changing the motion (acceleration) of charges or by changing magnetic fields. Electromagnetic waves can travel through matter, but they do not require a material medium. (That is, they also travel through empty space.) All electromagnetic waves move in a vacuum at the speed of light. Types of electromagnetic radiation are distinguished from each other by their wavelength and energy.

P4.6 Electromagnetic Waves

Electromagnetic waves (e.g., radio, microwave, infrared, visible light, ultraviolet, x-ray) are produced by changing the motion (acceleration) of charges or by changing magnetic fields. Electromagnetic waves can travel through matter, but they do not require a material medium. (That is, they also travel through empty space.) All electromagnetic waves move in a vacuum at the speed of light. Types of electromagnetic radiation are distinguished from each other by their wavelength and energy.

P4.6 Electromagnetic Waves

Electromagnetic waves (e.g., radio, microwave, infrared, visible light, ultraviolet, x-ray) are produced by changing the motion (acceleration) of charges or by changing magnetic fields. Electromagnetic waves can travel through matter, but they do not require a material medium. (That is, they also travel through empty space.) All electromagnetic waves move in a vacuum at the speed of light. Types of electromagnetic radiation are distinguished from each other by their wavelength and energy.

P4.6 Electromagnetic Waves

Electromagnetic waves (e.g., radio, microwave, infrared, visible light, ultraviolet, x-ray) are produced by changing the motion (acceleration) of charges or by changing magnetic fields. Electromagnetic waves can travel through matter, but they do not require a material medium. (That is, they also travel through empty space.) All electromagnetic waves move in a vacuum at the speed of light. Types of electromagnetic radiation are distinguished from each other by their wavelength and energy.

P4.6 Electromagnetic Waves

Electromagnetic waves (e.g., radio, microwave, infrared, visible light, ultraviolet, x-ray) are produced by changing the motion (acceleration) of charges or by changing magnetic fields. Electromagnetic waves can travel through matter, but they do not require a material medium. (That is, they also travel through empty space.) All electromagnetic waves move in a vacuum at the speed of light. Types of electromagnetic radiation are distinguished from each other by their wavelength and energy.

E

P4.6B Explain why radio waves can travel through space, but sound waves cannot.

E

P4.6B Explain why radio waves can travel through space, but sound waves cannot.

E

P4.6B Explain why radio waves can travel through space, but sound waves cannot.

E

P4.6B Explain why radio waves can travel through space, but sound waves cannot.

E

P4.6B Explain why radio waves can travel through space, but sound waves cannot.

E

P4.6B Explain why radio waves can travel through space, but sound waves cannot.

P4.6 Electromagnetic Waves

Electromagnetic waves (e.g., radio, microwave, infrared, visible light, ultraviolet, x-ray) are produced by changing the motion (acceleration) of charges or by changing magnetic fields. Electromagnetic waves can travel through matter, but they do not require a material medium. (That is, they also travel through empty space.) All electromagnetic waves move in a vacuum at the speed of light. Types of electromagnetic radiation are distinguished from each other by their wavelength and energy.

P4.6 Electromagnetic Waves

Electromagnetic waves (e.g., radio, microwave, infrared, visible light, ultraviolet, x-ray) are produced by changing the motion (acceleration) of charges or by changing magnetic fields. Electromagnetic waves can travel through matter, but they do not require a material medium. (That is, they also travel through empty space.) All electromagnetic waves move in a vacuum at the speed of light. Types of electromagnetic radiation are distinguished from each other by their wavelength and energy.

P4.6 Electromagnetic Waves

Electromagnetic waves (e.g., radio, microwave, infrared, visible light, ultraviolet, x-ray) are produced by changing the motion (acceleration) of charges or by changing magnetic fields. Electromagnetic waves can travel through matter, but they do not require a material medium. (That is, they also travel through empty space.) All electromagnetic waves move in a vacuum at the speed of light. Types of electromagnetic radiation are distinguished from each other by their wavelength and energy.

P4.6 Electromagnetic Waves

Electromagnetic waves (e.g., radio, microwave, infrared, visible light, ultraviolet, x-ray) are produced by changing the motion (acceleration) of charges or by changing magnetic fields. Electromagnetic waves can travel through matter, but they do not require a material medium. (That is, they also travel through empty space.) All electromagnetic waves move in a vacuum at the speed of light. Types of electromagnetic radiation are distinguished from each other by their wavelength and energy.

P4.6 Electromagnetic Waves

Electromagnetic waves (e.g., radio, microwave, infrared, visible light, ultraviolet, x-ray) are produced by changing the motion (acceleration) of charges or by changing magnetic fields. Electromagnetic waves can travel through matter, but they do not require a material medium. (That is, they also travel through empty space.) All electromagnetic waves move in a vacuum at the speed of light. Types of electromagnetic radiation are distinguished from each other by their wavelength and energy.

P4.6 Electromagnetic Waves

Electromagnetic waves (e.g., radio, microwave, infrared, visible light, ultraviolet, x-ray) are produced by changing the motion (acceleration) of charges or by changing magnetic fields. Electromagnetic waves can travel through matter, but they do not require a material medium. (That is, they also travel through empty space.) All electromagnetic waves move in a vacuum at the speed of light. Types of electromagnetic radiation are distinguished from each other by their wavelength and energy.

E

P4.6C Explain why there is a delay between the times we send a radio message to astronauts on the moon and when they receive it.

E

P4.6C Explain why there is a delay between the times we send a radio message to astronauts on the moon and when they receive it.

E

P4.6C Explain why there is a delay between the times we send a radio message to astronauts on the moon and when they receive it.

E

P4.6C Explain why there is a delay between the times we send a radio message to astronauts on the moon and when they receive it.

E

P4.6C Explain why there is a delay between the times we send a radio message to astronauts on the moon and when they receive it.

E

P4.6C Explain why there is a delay between the times we send a radio message to astronauts on the moon and when they receive it.

P4.6 Electromagnetic Waves

Electromagnetic waves (e.g., radio, microwave, infrared, visible light, ultraviolet, x-ray) are produced by changing the motion (acceleration) of charges or by changing magnetic fields. Electromagnetic waves can travel through matter, but they do not require a material medium. (That is, they also travel through empty space.) All electromagnetic waves move in a vacuum at the speed of light. Types of electromagnetic radiation are distinguished from each other by their wavelength and energy.

P4.6 Electromagnetic Waves

Electromagnetic waves (e.g., radio, microwave, infrared, visible light, ultraviolet, x-ray) are produced by changing the motion (acceleration) of charges or by changing magnetic fields. Electromagnetic waves can travel through matter, but they do not require a material medium. (That is, they also travel through empty space.) All electromagnetic waves move in a vacuum at the speed of light. Types of electromagnetic radiation are distinguished from each other by their wavelength and energy.

P4.6 Electromagnetic Waves

Electromagnetic waves (e.g., radio, microwave, infrared, visible light, ultraviolet, x-ray) are produced by changing the motion (acceleration) of charges or by changing magnetic fields. Electromagnetic waves can travel through matter, but they do not require a material medium. (That is, they also travel through empty space.) All electromagnetic waves move in a vacuum at the speed of light. Types of electromagnetic radiation are distinguished from each other by their wavelength and energy.

P4.6 Electromagnetic Waves

Electromagnetic waves (e.g., radio, microwave, infrared, visible light, ultraviolet, x-ray) are produced by changing the motion (acceleration) of charges or by changing magnetic fields. Electromagnetic waves can travel through matter, but they do not require a material medium. (That is, they also travel through empty space.) All electromagnetic waves move in a vacuum at the speed of light. Types of electromagnetic radiation are distinguished from each other by their wavelength and energy.

P4.6 Electromagnetic Waves

Electromagnetic waves (e.g., radio, microwave, infrared, visible light, ultraviolet, x-ray) are produced by changing the motion (acceleration) of charges or by changing magnetic fields. Electromagnetic waves can travel through matter, but they do not require a material medium. (That is, they also travel through empty space.) All electromagnetic waves move in a vacuum at the speed of light. Types of electromagnetic radiation are distinguished from each other by their wavelength and energy.

P4.6 Electromagnetic Waves

Electromagnetic waves (e.g., radio, microwave, infrared, visible light, ultraviolet, x-ray) are produced by changing the motion (acceleration) of charges or by changing magnetic fields. Electromagnetic waves can travel through matter, but they do not require a material medium. (That is, they also travel through empty space.) All electromagnetic waves move in a vacuum at the speed of light. Types of electromagnetic radiation are distinguished from each other by their wavelength and energy.

E

P4.6D Explain why we see a distant event before we hear it (e.g., lightning before thunder, exploding fireworks before the boom).

E

P4.6D Explain why we see a distant event before we hear it (e.g., lightning before thunder, exploding fireworks before the boom).

E

P4.6D Explain why we see a distant event before we hear it (e.g., lightning before thunder, exploding fireworks before the boom).

E

P4.6D Explain why we see a distant event before we hear it (e.g., lightning before thunder, exploding fireworks before the boom).

E

P4.6D Explain why we see a distant event before we hear it (e.g., lightning before thunder, exploding fireworks before the boom).

E

P4.6D Explain why we see a distant event before we hear it (e.g., lightning before thunder, exploding fireworks before the boom).

P4.6x Electromagnetic Propagation

Modulated electromagnetic waves can transfer information from one place to another (e.g., televisions, radios, telephones, computers and other information technology devices). Digital communication makes more efficient use of the limited electromagnetic spectrum, is more accurate than analog transmission, and can be encrypted to provide privacy and security.

P4.6x Electromagnetic Propagation

Modulated electromagnetic waves can transfer information from one place to another (e.g., televisions, radios, telephones, computers and other information technology devices). Digital communication makes more efficient use of the limited electromagnetic spectrum, is more accurate than analog transmission, and can be encrypted to provide privacy and security.

P4.6x Electromagnetic Propagation

Modulated electromagnetic waves can transfer information from one place to another (e.g., televisions, radios, telephones, computers and other information technology devices). Digital communication makes more efficient use of the limited electromagnetic spectrum, is more accurate than analog transmission, and can be encrypted to provide privacy and security.

P4.6x Electromagnetic Propagation

Modulated electromagnetic waves can transfer information from one place to another (e.g., televisions, radios, telephones, computers and other information technology devices). Digital communication makes more efficient use of the limited electromagnetic spectrum, is more accurate than analog transmission, and can be encrypted to provide privacy and security.

P4.6x Electromagnetic Propagation

Modulated electromagnetic waves can transfer information from one place to another (e.g., televisions, radios, telephones, computers and other information technology devices). Digital communication makes more efficient use of the limited electromagnetic spectrum, is more accurate than analog transmission, and can be encrypted to provide privacy and security.

P4.6x Electromagnetic Propagation

Modulated electromagnetic waves can transfer information from one place to another (e.g., televisions, radios, telephones, computers and other information technology devices). Digital communication makes more efficient use of the limited electromagnetic spectrum, is more accurate than analog transmission, and can be encrypted to provide privacy and security.



P4.6e Explain why antennas are needed for radio, television, and cell phone transmission and reception.



P4.6e Explain why antennas are needed for radio, television, and cell phone transmission and reception.



P4.6e Explain why antennas are needed for radio, television, and cell phone transmission and reception.



P4.6e Explain why antennas are needed for radio, television, and cell phone transmission and reception.



P4.6e Explain why antennas are needed for radio, television, and cell phone transmission and reception.



P4.6e Explain why antennas are needed for radio, television, and cell phone transmission and reception.

P4.6x Electromagnetic Propagation

Modulated electromagnetic waves can transfer information from one place to another (e.g., televisions, radios, telephones, computers and other information technology devices). Digital communication makes more efficient use of the limited electromagnetic spectrum, is more accurate than analog transmission, and can be encrypted to provide privacy and security.

P4.6x Electromagnetic Propagation

Modulated electromagnetic waves can transfer information from one place to another (e.g., televisions, radios, telephones, computers and other information technology devices). Digital communication makes more efficient use of the limited electromagnetic spectrum, is more accurate than analog transmission, and can be encrypted to provide privacy and security.

P4.6x Electromagnetic Propagation

Modulated electromagnetic waves can transfer information from one place to another (e.g., televisions, radios, telephones, computers and other information technology devices). Digital communication makes more efficient use of the limited electromagnetic spectrum, is more accurate than analog transmission, and can be encrypted to provide privacy and security.

P4.6x Electromagnetic Propagation

Modulated electromagnetic waves can transfer information from one place to another (e.g., televisions, radios, telephones, computers and other information technology devices). Digital communication makes more efficient use of the limited electromagnetic spectrum, is more accurate than analog transmission, and can be encrypted to provide privacy and security.

P4.6x Electromagnetic Propagation

Modulated electromagnetic waves can transfer information from one place to another (e.g., televisions, radios, telephones, computers and other information technology devices). Digital communication makes more efficient use of the limited electromagnetic spectrum, is more accurate than analog transmission, and can be encrypted to provide privacy and security.

P4.6x Electromagnetic Propagation

Modulated electromagnetic waves can transfer information from one place to another (e.g., televisions, radios, telephones, computers and other information technology devices). Digital communication makes more efficient use of the limited electromagnetic spectrum, is more accurate than analog transmission, and can be encrypted to provide privacy and security.



P4.6f Explain how radio waves are modified to send information in radio and television programs, radio-controlled cars, cell phone conversations, and GPS systems.



P4.6f Explain how radio waves are modified to send information in radio and television programs, radio-controlled cars, cell phone conversations, and GPS systems.



P4.6f Explain how radio waves are modified to send information in radio and television programs, radio-controlled cars, cell phone conversations, and GPS systems.



P4.6f Explain how radio waves are modified to send information in radio and television programs, radio-controlled cars, cell phone conversations, and GPS systems.



P4.6f Explain how radio waves are modified to send information in radio and television programs, radio-controlled cars, cell phone conversations, and GPS systems.



P4.6f Explain how radio waves are modified to send information in radio and television programs, radio-controlled cars, cell phone conversations, and GPS systems.

P4.6x Electromagnetic Propagation

Modulated electromagnetic waves can transfer information from one place to another (e.g., televisions, radios, telephones, computers and other information technology devices). Digital communication makes more efficient use of the limited electromagnetic spectrum, is more accurate than analog transmission, and can be encrypted to provide privacy and security.

P4.6x Electromagnetic Propagation

Modulated electromagnetic waves can transfer information from one place to another (e.g., televisions, radios, telephones, computers and other information technology devices). Digital communication makes more efficient use of the limited electromagnetic spectrum, is more accurate than analog transmission, and can be encrypted to provide privacy and security.

P4.6x Electromagnetic Propagation

Modulated electromagnetic waves can transfer information from one place to another (e.g., televisions, radios, telephones, computers and other information technology devices). Digital communication makes more efficient use of the limited electromagnetic spectrum, is more accurate than analog transmission, and can be encrypted to provide privacy and security.

P4.6x Electromagnetic Propagation

Modulated electromagnetic waves can transfer information from one place to another (e.g., televisions, radios, telephones, computers and other information technology devices). Digital communication makes more efficient use of the limited electromagnetic spectrum, is more accurate than analog transmission, and can be encrypted to provide privacy and security.

P4.6x Electromagnetic Propagation

Modulated electromagnetic waves can transfer information from one place to another (e.g., televisions, radios, telephones, computers and other information technology devices). Digital communication makes more efficient use of the limited electromagnetic spectrum, is more accurate than analog transmission, and can be encrypted to provide privacy and security.

P4.6x Electromagnetic Propagation

Modulated electromagnetic waves can transfer information from one place to another (e.g., televisions, radios, telephones, computers and other information technology devices). Digital communication makes more efficient use of the limited electromagnetic spectrum, is more accurate than analog transmission, and can be encrypted to provide privacy and security.



P4.6g Explain how different electromagnetic signals (e.g., radio station broadcasts or cell phone conversations) can take place without interfering with each other.



P4.6g Explain how different electromagnetic signals (e.g., radio station broadcasts or cell phone conversations) can take place without interfering with each other.



P4.6g Explain how different electromagnetic signals (e.g., radio station broadcasts or cell phone conversations) can take place without interfering with each other.



P4.6g Explain how different electromagnetic signals (e.g., radio station broadcasts or cell phone conversations) can take place without interfering with each other.



P4.6g Explain how different electromagnetic signals (e.g., radio station broadcasts or cell phone conversations) can take place without interfering with each other.



P4.6g Explain how different electromagnetic signals (e.g., radio station broadcasts or cell phone conversations) can take place without interfering with each other.

P4.6x Electromagnetic Propagation

Modulated electromagnetic waves can transfer information from one place to another (e.g., televisions, radios, telephones, computers and other information technology devices). Digital communication makes more efficient use of the limited electromagnetic spectrum, is more accurate than analog transmission, and can be encrypted to provide privacy and security.

P4.6x Electromagnetic Propagation

Modulated electromagnetic waves can transfer information from one place to another (e.g., televisions, radios, telephones, computers and other information technology devices). Digital communication makes more efficient use of the limited electromagnetic spectrum, is more accurate than analog transmission, and can be encrypted to provide privacy and security.

P4.6x Electromagnetic Propagation

Modulated electromagnetic waves can transfer information from one place to another (e.g., televisions, radios, telephones, computers and other information technology devices). Digital communication makes more efficient use of the limited electromagnetic spectrum, is more accurate than analog transmission, and can be encrypted to provide privacy and security.

P4.6x Electromagnetic Propagation

Modulated electromagnetic waves can transfer information from one place to another (e.g., televisions, radios, telephones, computers and other information technology devices). Digital communication makes more efficient use of the limited electromagnetic spectrum, is more accurate than analog transmission, and can be encrypted to provide privacy and security.

P4.6x Electromagnetic Propagation

Modulated electromagnetic waves can transfer information from one place to another (e.g., televisions, radios, telephones, computers and other information technology devices). Digital communication makes more efficient use of the limited electromagnetic spectrum, is more accurate than analog transmission, and can be encrypted to provide privacy and security.

P4.6x Electromagnetic Propagation

Modulated electromagnetic waves can transfer information from one place to another (e.g., televisions, radios, telephones, computers and other information technology devices). Digital communication makes more efficient use of the limited electromagnetic spectrum, is more accurate than analog transmission, and can be encrypted to provide privacy and security.



P4.6h Explain the relationship between the frequency of an electromagnetic wave and its technological uses.



P4.6h Explain the relationship between the frequency of an electromagnetic wave and its technological uses.



P4.6h Explain the relationship between the frequency of an electromagnetic wave and its technological uses.



P4.6h Explain the relationship between the frequency of an electromagnetic wave and its technological uses.



P4.6h Explain the relationship between the frequency of an electromagnetic wave and its technological uses.



P4.6h Explain the relationship between the frequency of an electromagnetic wave and its technological uses.

P4.r7x Quantum Theory of Waves (recommended)

Electromagnetic energy is transferred on the atomic scale in discrete amounts called quanta. The equation $E = hf$ quantifies the relationship between the energy transferred and the frequency, where h is Planck's constant. (recommended)

P4.r7x Quantum Theory of Waves (recommended)

Electromagnetic energy is transferred on the atomic scale in discrete amounts called quanta. The equation $E = hf$ quantifies the relationship between the energy transferred and the frequency, where h is Planck's constant. (recommended)

P4.r7x Quantum Theory of Waves (recommended)

Electromagnetic energy is transferred on the atomic scale in discrete amounts called quanta. The equation $E = hf$ quantifies the relationship between the energy transferred and the frequency, where h is Planck's constant. (recommended)

P4.r7x Quantum Theory of Waves (recommended)

Electromagnetic energy is transferred on the atomic scale in discrete amounts called quanta. The equation $E = hf$ quantifies the relationship between the energy transferred and the frequency, where h is Planck's constant. (recommended)

P4.r7x Quantum Theory of Waves (recommended)

Electromagnetic energy is transferred on the atomic scale in discrete amounts called quanta. The equation $E = hf$ quantifies the relationship between the energy transferred and the frequency, where h is Planck's constant. (recommended)

P4.r7x Quantum Theory of Waves (recommended)

Electromagnetic energy is transferred on the atomic scale in discrete amounts called quanta. The equation $E = hf$ quantifies the relationship between the energy transferred and the frequency, where h is Planck's constant. (recommended)



P4.r7h Calculate and compare the energy in various electromagnetic quanta (e.g., visible light, x-rays).(recommended)



P4.r7h Calculate and compare the energy in various electromagnetic quanta (e.g., visible light, x-rays).(recommended)



P4.r7h Calculate and compare the energy in various electromagnetic quanta (e.g., visible light, x-rays).(recommended)



P4.r7h Calculate and compare the energy in various electromagnetic quanta (e.g., visible light, x-rays).(recommended)



P4.r7h Calculate and compare the energy in various electromagnetic quanta (e.g., visible light, x-rays).(recommended)



P4.r7h Calculate and compare the energy in various electromagnetic quanta (e.g., visible light, x-rays).(recommended)

P4.8 Wave Behavior — Reflection and Refraction

The laws of reflection and refraction describe the relationships between incident and reflected/refracted waves.

P4.8 Wave Behavior — Reflection and Refraction

The laws of reflection and refraction describe the relationships between incident and reflected/refracted waves.

P4.8 Wave Behavior — Reflection and Refraction

The laws of reflection and refraction describe the relationships between incident and reflected/refracted waves.

P4.8 Wave Behavior — Reflection and Refraction

The laws of reflection and refraction describe the relationships between incident and reflected/refracted waves.

P4.8 Wave Behavior — Reflection and Refraction

The laws of reflection and refraction describe the relationships between incident and reflected/refracted waves.

P4.8 Wave Behavior — Reflection and Refraction

The laws of reflection and refraction describe the relationships between incident and reflected/refracted waves.



P4.8A Draw ray diagrams to indicate how light reflects off objects or refracts into transparent media.



P4.8A Draw ray diagrams to indicate how light reflects off objects or refracts into transparent media.



P4.8A Draw ray diagrams to indicate how light reflects off objects or refracts into transparent media.



P4.8A Draw ray diagrams to indicate how light reflects off objects or refracts into transparent media.



P4.8A Draw ray diagrams to indicate how light reflects off objects or refracts into transparent media.



P4.8A Draw ray diagrams to indicate how light reflects off objects or refracts into transparent media.

P4.8 Wave Behavior — Reflection and Refraction

The laws of reflection and refraction describe the relationships between incident and reflected/refracted waves.

P4.8 Wave Behavior — Reflection and Refraction

The laws of reflection and refraction describe the relationships between incident and reflected/refracted waves.

P4.8 Wave Behavior — Reflection and Refraction

The laws of reflection and refraction describe the relationships between incident and reflected/refracted waves.

P4.8 Wave Behavior — Reflection and Refraction

The laws of reflection and refraction describe the relationships between incident and reflected/refracted waves.

P4.8 Wave Behavior — Reflection and Refraction

The laws of reflection and refraction describe the relationships between incident and reflected/refracted waves.

P4.8 Wave Behavior — Reflection and Refraction

The laws of reflection and refraction describe the relationships between incident and reflected/refracted waves.



P4.8B Predict the path of reflected light from flat, curved, or rough surfaces (e.g., flat and curved mirrors, painted walls, paper).



P4.8B Predict the path of reflected light from flat, curved, or rough surfaces (e.g., flat and curved mirrors, painted walls, paper).



P4.8B Predict the path of reflected light from flat, curved, or rough surfaces (e.g., flat and curved mirrors, painted walls, paper).



P4.8B Predict the path of reflected light from flat, curved, or rough surfaces (e.g., flat and curved mirrors, painted walls, paper).



P4.8B Predict the path of reflected light from flat, curved, or rough surfaces (e.g., flat and curved mirrors, painted walls, paper).



P4.8B Predict the path of reflected light from flat, curved, or rough surfaces (e.g., flat and curved mirrors, painted walls, paper).

P4.8x Wave Behavior — Diffraction, Interference, and Refraction

Waves can bend around objects (diffraction). They also superimpose on each other and continue their propagation without a change in their original properties (interference). When refracted, light follows a defined path.

P4.8x Wave Behavior — Diffraction, Interference, and Refraction

Waves can bend around objects (diffraction). They also superimpose on each other and continue their propagation without a change in their original properties (interference). When refracted, light follows a defined path.

P4.8x Wave Behavior — Diffraction, Interference, and Refraction

Waves can bend around objects (diffraction). They also superimpose on each other and continue their propagation without a change in their original properties (interference). When refracted, light follows a defined path.

P4.8x Wave Behavior — Diffraction, Interference, and Refraction

Waves can bend around objects (diffraction). They also superimpose on each other and continue their propagation without a change in their original properties (interference). When refracted, light follows a defined path.

P4.8x Wave Behavior — Diffraction, Interference, and Refraction

Waves can bend around objects (diffraction). They also superimpose on each other and continue their propagation without a change in their original properties (interference). When refracted, light follows a defined path.

P4.8x Wave Behavior — Diffraction, Interference, and Refraction

Waves can bend around objects (diffraction). They also superimpose on each other and continue their propagation without a change in their original properties (interference). When refracted, light follows a defined path.



P4.8c Describe how two wave pulses propagated from opposite ends of a demonstration spring interact as they meet.



P4.8c Describe how two wave pulses propagated from opposite ends of a demonstration spring interact as they meet.



P4.8c Describe how two wave pulses propagated from opposite ends of a demonstration spring interact as they meet.



P4.8c Describe how two wave pulses propagated from opposite ends of a demonstration spring interact as they meet.



P4.8c Describe how two wave pulses propagated from opposite ends of a demonstration spring interact as they meet.



P4.8c Describe how two wave pulses propagated from opposite ends of a demonstration spring interact as they meet.

P4.8x Wave Behavior — Diffraction, Interference, and Refraction

Waves can bend around objects (diffraction). They also superimpose on each other and continue their propagation without a change in their original properties (interference). When refracted, light follows a defined path.

P4.8x Wave Behavior — Diffraction, Interference, and Refraction

Waves can bend around objects (diffraction). They also superimpose on each other and continue their propagation without a change in their original properties (interference). When refracted, light follows a defined path.

P4.8x Wave Behavior — Diffraction, Interference, and Refraction

Waves can bend around objects (diffraction). They also superimpose on each other and continue their propagation without a change in their original properties (interference). When refracted, light follows a defined path.

P4.8x Wave Behavior — Diffraction, Interference, and Refraction

Waves can bend around objects (diffraction). They also superimpose on each other and continue their propagation without a change in their original properties (interference). When refracted, light follows a defined path.

P4.8x Wave Behavior — Diffraction, Interference, and Refraction

Waves can bend around objects (diffraction). They also superimpose on each other and continue their propagation without a change in their original properties (interference). When refracted, light follows a defined path.

P4.8x Wave Behavior — Diffraction, Interference, and Refraction

Waves can bend around objects (diffraction). They also superimpose on each other and continue their propagation without a change in their original properties (interference). When refracted, light follows a defined path.



P4.8d List and analyze everyday examples that demonstrate the interference characteristics of waves (e.g., dead spots in an auditorium, whispering galleries, colors in a CD, beetle wings).



P4.8d List and analyze everyday examples that demonstrate the interference characteristics of waves (e.g., dead spots in an auditorium, whispering galleries, colors in a CD, beetle wings).



P4.8d List and analyze everyday examples that demonstrate the interference characteristics of waves (e.g., dead spots in an auditorium, whispering galleries, colors in a CD, beetle wings).



P4.8d List and analyze everyday examples that demonstrate the interference characteristics of waves (e.g., dead spots in an auditorium, whispering galleries, colors in a CD, beetle wings).



P4.8d List and analyze everyday examples that demonstrate the interference characteristics of waves (e.g., dead spots in an auditorium, whispering galleries, colors in a CD, beetle wings).



P4.8d List and analyze everyday examples that demonstrate the interference characteristics of waves (e.g., dead spots in an auditorium, whispering galleries, colors in a CD, beetle wings).

P4.8x Wave Behavior — Diffraction, Interference, and Refraction

Waves can bend around objects (diffraction). They also superimpose on each other and continue their propagation without a change in their original properties (interference). When refracted, light follows a defined path.

P4.8x Wave Behavior — Diffraction, Interference, and Refraction

Waves can bend around objects (diffraction). They also superimpose on each other and continue their propagation without a change in their original properties (interference). When refracted, light follows a defined path.

P4.8x Wave Behavior — Diffraction, Interference, and Refraction

Waves can bend around objects (diffraction). They also superimpose on each other and continue their propagation without a change in their original properties (interference). When refracted, light follows a defined path.

P4.8x Wave Behavior — Diffraction, Interference, and Refraction

Waves can bend around objects (diffraction). They also superimpose on each other and continue their propagation without a change in their original properties (interference). When refracted, light follows a defined path.

P4.8x Wave Behavior — Diffraction, Interference, and Refraction

Waves can bend around objects (diffraction). They also superimpose on each other and continue their propagation without a change in their original properties (interference). When refracted, light follows a defined path.

P4.8x Wave Behavior — Diffraction, Interference, and Refraction

Waves can bend around objects (diffraction). They also superimpose on each other and continue their propagation without a change in their original properties (interference). When refracted, light follows a defined path.



P4.8e Given an angle of incidence and indices of refraction of two materials, calculate the path of a light ray incident on the boundary (Snell's Law).



P4.8e Given an angle of incidence and indices of refraction of two materials, calculate the path of a light ray incident on the boundary (Snell's Law).



P4.8e Given an angle of incidence and indices of refraction of two materials, calculate the path of a light ray incident on the boundary (Snell's Law).



P4.8e Given an angle of incidence and indices of refraction of two materials, calculate the path of a light ray incident on the boundary (Snell's Law).



P4.8e Given an angle of incidence and indices of refraction of two materials, calculate the path of a light ray incident on the boundary (Snell's Law).



P4.8e Given an angle of incidence and indices of refraction of two materials, calculate the path of a light ray incident on the boundary (Snell's Law).

P4.8x Wave Behavior — Diffraction, Interference, and Refraction

Waves can bend around objects (diffraction). They also superimpose on each other and continue their propagation without a change in their original properties (interference). When refracted, light follows a defined path.

P4.8x Wave Behavior — Diffraction, Interference, and Refraction

Waves can bend around objects (diffraction). They also superimpose on each other and continue their propagation without a change in their original properties (interference). When refracted, light follows a defined path.

P4.8x Wave Behavior — Diffraction, Interference, and Refraction

Waves can bend around objects (diffraction). They also superimpose on each other and continue their propagation without a change in their original properties (interference). When refracted, light follows a defined path.

P4.8x Wave Behavior — Diffraction, Interference, and Refraction

Waves can bend around objects (diffraction). They also superimpose on each other and continue their propagation without a change in their original properties (interference). When refracted, light follows a defined path.

P4.8x Wave Behavior — Diffraction, Interference, and Refraction

Waves can bend around objects (diffraction). They also superimpose on each other and continue their propagation without a change in their original properties (interference). When refracted, light follows a defined path.

P4.8x Wave Behavior — Diffraction, Interference, and Refraction

Waves can bend around objects (diffraction). They also superimpose on each other and continue their propagation without a change in their original properties (interference). When refracted, light follows a defined path.



P4.8f Explain how Snell's Law is used to design lenses (e.g., eye glasses, microscopes, telescopes, binoculars).



P4.8f Explain how Snell's Law is used to design lenses (e.g., eye glasses, microscopes, telescopes, binoculars).



P4.8f Explain how Snell's Law is used to design lenses (e.g., eye glasses, microscopes, telescopes, binoculars).



P4.8f Explain how Snell's Law is used to design lenses (e.g., eye glasses, microscopes, telescopes, binoculars).



P4.8f Explain how Snell's Law is used to design lenses (e.g., eye glasses, microscopes, telescopes, binoculars).



P4.8f Explain how Snell's Law is used to design lenses (e.g., eye glasses, microscopes, telescopes, binoculars).

P4.9 Nature of Light

Light interacts with matter by reflection, absorption, or transmission.

P4.9 Nature of Light

Light interacts with matter by reflection, absorption, or transmission.

P4.9 Nature of Light

Light interacts with matter by reflection, absorption, or transmission.

P4.9 Nature of Light

Light interacts with matter by reflection, absorption, or transmission.

P4.9 Nature of Light

Light interacts with matter by reflection, absorption, or transmission.

P4.9 Nature of Light

Light interacts with matter by reflection, absorption, or transmission.



P4.9A Identify the principle involved when you see a transparent object (e.g., straw, piece of glass) in a clear liquid.



P4.9A Identify the principle involved when you see a transparent object (e.g., straw, piece of glass) in a clear liquid.



P4.9A Identify the principle involved when you see a transparent object (e.g., straw, piece of glass) in a clear liquid.



P4.9A Identify the principle involved when you see a transparent object (e.g., straw, piece of glass) in a clear liquid.



P4.9A Identify the principle involved when you see a transparent object (e.g., straw, piece of glass) in a clear liquid.



P4.9A Identify the principle involved when you see a transparent object (e.g., straw, piece of glass) in a clear liquid.

P4.9 Nature of Light

Light interacts with matter by reflection, absorption, or transmission.

P4.9 Nature of Light

Light interacts with matter by reflection, absorption, or transmission.

P4.9 Nature of Light

Light interacts with matter by reflection, absorption, or transmission.

P4.9 Nature of Light

Light interacts with matter by reflection, absorption, or transmission.

P4.9 Nature of Light

Light interacts with matter by reflection, absorption, or transmission.

P4.9 Nature of Light

Light interacts with matter by reflection, absorption, or transmission.



P4.9B Explain how various materials reflect, absorb, or transmit light in different ways.



P4.9B Explain how various materials reflect, absorb, or transmit light in different ways.



P4.9B Explain how various materials reflect, absorb, or transmit light in different ways.



P4.9B Explain how various materials reflect, absorb, or transmit light in different ways.



P4.9B Explain how various materials reflect, absorb, or transmit light in different ways.



P4.9B Explain how various materials reflect, absorb, or transmit light in different ways.

P4.9 Nature of Light

Light interacts with matter by reflection, absorption, or transmission.

P4.9 Nature of Light

Light interacts with matter by reflection, absorption, or transmission.

P4.9 Nature of Light

Light interacts with matter by reflection, absorption, or transmission.

P4.9 Nature of Light

Light interacts with matter by reflection, absorption, or transmission.

P4.9 Nature of Light

Light interacts with matter by reflection, absorption, or transmission.

P4.9 Nature of Light

Light interacts with matter by reflection, absorption, or transmission.



P4.9C Explain why the image of the Sun appears red-dish at sunrise and sunset.



P4.9C Explain why the image of the Sun appears red-dish at sunrise and sunset.



P4.9C Explain why the image of the Sun appears red-dish at sunrise and sunset.



P4.9C Explain why the image of the Sun appears red-dish at sunrise and sunset.



P4.9C Explain why the image of the Sun appears red-dish at sunrise and sunset.



P4.9C Explain why the image of the Sun appears red-dish at sunrise and sunset.

P4.r9x Nature of Light —

Wave-Particle Nature (recommended)

The dual wave-particle nature of matter and light is the foundation for modern physics. (recommended)

P4.r9x Nature of Light —

Wave-Particle Nature (recommended)

The dual wave-particle nature of matter and light is the foundation for modern physics. (recommended)

P4.r9x Nature of Light —

Wave-Particle Nature (recommended)

The dual wave-particle nature of matter and light is the foundation for modern physics. (recommended)

P4.r9x Nature of Light —

Wave-Particle Nature (recommended)

The dual wave-particle nature of matter and light is the foundation for modern physics. (recommended)

P4.r9x Nature of Light —

Wave-Particle Nature (recommended)

The dual wave-particle nature of matter and light is the foundation for modern physics. (recommended)

P4.r9x Nature of Light —

Wave-Particle Nature (recommended)

The dual wave-particle nature of matter and light is the foundation for modern physics. (recommended)



P4.r9d Describe evidence that supports the dual wave - particle nature of light. (recommended)



P4.r9d Describe evidence that supports the dual wave - particle nature of light. (recommended)



P4.r9d Describe evidence that supports the dual wave - particle nature of light. (recommended)



P4.r9d Describe evidence that supports the dual wave - particle nature of light. (recommended)



P4.r9d Describe evidence that supports the dual wave - particle nature of light. (recommended)



P4.r9d Describe evidence that supports the dual wave - particle nature of light. (recommended)

P4.10 Current Electricity — Circuits

Current electricity is described as movement of charges. It is a particularly useful form of energy because it can be easily transferred from place to place and readily transformed by various devices into other forms of energy (e.g., light, heat, sound, and motion). Electrical current (amperage) in a circuit is determined by the potential difference (voltage) of the power source and the resistance of the loads in the circuit.

P4.10 Current Electricity — Circuits

Current electricity is described as movement of charges. It is a particularly useful form of energy because it can be easily transferred from place to place and readily transformed by various devices into other forms of energy (e.g., light, heat, sound, and motion). Electrical current (amperage) in a circuit is determined by the potential difference (voltage) of the power source and the resistance of the loads in the circuit.

P4.10 Current Electricity — Circuits

Current electricity is described as movement of charges. It is a particularly useful form of energy because it can be easily transferred from place to place and readily transformed by various devices into other forms of energy (e.g., light, heat, sound, and motion). Electrical current (amperage) in a circuit is determined by the potential difference (voltage) of the power source and the resistance of the loads in the circuit.

P4.10 Current Electricity — Circuits

Current electricity is described as movement of charges. It is a particularly useful form of energy because it can be easily transferred from place to place and readily transformed by various devices into other forms of energy (e.g., light, heat, sound, and motion). Electrical current (amperage) in a circuit is determined by the potential difference (voltage) of the power source and the resistance of the loads in the circuit.

P4.10 Current Electricity — Circuits

Current electricity is described as movement of charges. It is a particularly useful form of energy because it can be easily transferred from place to place and readily transformed by various devices into other forms of energy (e.g., light, heat, sound, and motion). Electrical current (amperage) in a circuit is determined by the potential difference (voltage) of the power source and the resistance of the loads in the circuit.

P4.10 Current Electricity — Circuits

Current electricity is described as movement of charges. It is a particularly useful form of energy because it can be easily transferred from place to place and readily transformed by various devices into other forms of energy (e.g., light, heat, sound, and motion). Electrical current (amperage) in a circuit is determined by the potential difference (voltage) of the power source and the resistance of the loads in the circuit.



P4.10A Describe the energy transformations when electrical energy is produced and transferred to homes and businesses.



P4.10A Describe the energy transformations when electrical energy is produced and transferred to homes and businesses.



P4.10A Describe the energy transformations when electrical energy is produced and transferred to homes and businesses.



P4.10A Describe the energy transformations when electrical energy is produced and transferred to homes and businesses.



P4.10A Describe the energy transformations when electrical energy is produced and transferred to homes and businesses.



P4.10A Describe the energy transformations when electrical energy is produced and transferred to homes and businesses.

P4.10 Current Electricity — Circuits

Current electricity is described as movement of charges. It is a particularly useful form of energy because it can be easily transferred from place to place and readily transformed by various devices into other forms of energy (e.g., light, heat, sound, and motion). Electrical current (amperage) in a circuit is determined by the potential difference (voltage) of the power source and the resistance of the loads in the circuit.

P4.10 Current Electricity — Circuits

Current electricity is described as movement of charges. It is a particularly useful form of energy because it can be easily transferred from place to place and readily transformed by various devices into other forms of energy (e.g., light, heat, sound, and motion). Electrical current (amperage) in a circuit is determined by the potential difference (voltage) of the power source and the resistance of the loads in the circuit.

P4.10 Current Electricity — Circuits

Current electricity is described as movement of charges. It is a particularly useful form of energy because it can be easily transferred from place to place and readily transformed by various devices into other forms of energy (e.g., light, heat, sound, and motion). Electrical current (amperage) in a circuit is determined by the potential difference (voltage) of the power source and the resistance of the loads in the circuit.

P4.10 Current Electricity — Circuits

Current electricity is described as movement of charges. It is a particularly useful form of energy because it can be easily transferred from place to place and readily transformed by various devices into other forms of energy (e.g., light, heat, sound, and motion). Electrical current (amperage) in a circuit is determined by the potential difference (voltage) of the power source and the resistance of the loads in the circuit.

P4.10 Current Electricity — Circuits

Current electricity is described as movement of charges. It is a particularly useful form of energy because it can be easily transferred from place to place and readily transformed by various devices into other forms of energy (e.g., light, heat, sound, and motion). Electrical current (amperage) in a circuit is determined by the potential difference (voltage) of the power source and the resistance of the loads in the circuit.

P4.10 Current Electricity — Circuits

Current electricity is described as movement of charges. It is a particularly useful form of energy because it can be easily transferred from place to place and readily transformed by various devices into other forms of energy (e.g., light, heat, sound, and motion). Electrical current (amperage) in a circuit is determined by the potential difference (voltage) of the power source and the resistance of the loads in the circuit.



P4.10B Identify common household devices that transform electrical energy to other forms of energy, and describe the type of energy transformation.



P4.10B Identify common household devices that transform electrical energy to other forms of energy, and describe the type of energy transformation.



P4.10B Identify common household devices that transform electrical energy to other forms of energy, and describe the type of energy transformation.



P4.10B Identify common household devices that transform electrical energy to other forms of energy, and describe the type of energy transformation.



P4.10B Identify common household devices that transform electrical energy to other forms of energy, and describe the type of energy transformation.



P4.10B Identify common household devices that transform electrical energy to other forms of energy, and describe the type of energy transformation.

P4.10 Current Electricity — Circuits

Current electricity is described as movement of charges. It is a particularly useful form of energy because it can be easily transferred from place to place and readily transformed by various devices into other forms of energy (e.g., light, heat, sound, and motion). Electrical current (amperage) in a circuit is determined by the potential difference (voltage) of the power source and the resistance of the loads in the circuit.

P4.10 Current Electricity — Circuits

Current electricity is described as movement of charges. It is a particularly useful form of energy because it can be easily transferred from place to place and readily transformed by various devices into other forms of energy (e.g., light, heat, sound, and motion). Electrical current (amperage) in a circuit is determined by the potential difference (voltage) of the power source and the resistance of the loads in the circuit.

P4.10 Current Electricity — Circuits

Current electricity is described as movement of charges. It is a particularly useful form of energy because it can be easily transferred from place to place and readily transformed by various devices into other forms of energy (e.g., light, heat, sound, and motion). Electrical current (amperage) in a circuit is determined by the potential difference (voltage) of the power source and the resistance of the loads in the circuit.

P4.10 Current Electricity — Circuits

Current electricity is described as movement of charges. It is a particularly useful form of energy because it can be easily transferred from place to place and readily transformed by various devices into other forms of energy (e.g., light, heat, sound, and motion). Electrical current (amperage) in a circuit is determined by the potential difference (voltage) of the power source and the resistance of the loads in the circuit.

P4.10 Current Electricity — Circuits

Current electricity is described as movement of charges. It is a particularly useful form of energy because it can be easily transferred from place to place and readily transformed by various devices into other forms of energy (e.g., light, heat, sound, and motion). Electrical current (amperage) in a circuit is determined by the potential difference (voltage) of the power source and the resistance of the loads in the circuit.

P4.10 Current Electricity — Circuits

Current electricity is described as movement of charges. It is a particularly useful form of energy because it can be easily transferred from place to place and readily transformed by various devices into other forms of energy (e.g., light, heat, sound, and motion). Electrical current (amperage) in a circuit is determined by the potential difference (voltage) of the power source and the resistance of the loads in the circuit.



P4.10C Given diagrams of many different possible connections of electric circuit elements, identify complete circuits, open circuits, and short circuits and explain the reasons for the classification.



P4.10C Given diagrams of many different possible connections of electric circuit elements, identify complete circuits, open circuits, and short circuits and explain the reasons for the classification.



P4.10C Given diagrams of many different possible connections of electric circuit elements, identify complete circuits, open circuits, and short circuits and explain the reasons for the classification.



P4.10C Given diagrams of many different possible connections of electric circuit elements, identify complete circuits, open circuits, and short circuits and explain the reasons for the classification.



P4.10C Given diagrams of many different possible connections of electric circuit elements, identify complete circuits, open circuits, and short circuits and explain the reasons for the classification.



P4.10C Given diagrams of many different possible connections of electric circuit elements, identify complete circuits, open circuits, and short circuits and explain the reasons for the classification.

P4.10 Current Electricity — Circuits

Current electricity is described as movement of charges. It is a particularly useful form of energy because it can be easily transferred from place to place and readily transformed by various devices into other forms of energy (e.g., light, heat, sound, and motion). Electrical current (amperage) in a circuit is determined by the potential difference (voltage) of the power source and the resistance of the loads in the circuit.

P4.10 Current Electricity — Circuits

Current electricity is described as movement of charges. It is a particularly useful form of energy because it can be easily transferred from place to place and readily transformed by various devices into other forms of energy (e.g., light, heat, sound, and motion). Electrical current (amperage) in a circuit is determined by the potential difference (voltage) of the power source and the resistance of the loads in the circuit.

P4.10 Current Electricity — Circuits

Current electricity is described as movement of charges. It is a particularly useful form of energy because it can be easily transferred from place to place and readily transformed by various devices into other forms of energy (e.g., light, heat, sound, and motion). Electrical current (amperage) in a circuit is determined by the potential difference (voltage) of the power source and the resistance of the loads in the circuit.

P4.10 Current Electricity — Circuits

Current electricity is described as movement of charges. It is a particularly useful form of energy because it can be easily transferred from place to place and readily transformed by various devices into other forms of energy (e.g., light, heat, sound, and motion). Electrical current (amperage) in a circuit is determined by the potential difference (voltage) of the power source and the resistance of the loads in the circuit.

P4.10 Current Electricity — Circuits

Current electricity is described as movement of charges. It is a particularly useful form of energy because it can be easily transferred from place to place and readily transformed by various devices into other forms of energy (e.g., light, heat, sound, and motion). Electrical current (amperage) in a circuit is determined by the potential difference (voltage) of the power source and the resistance of the loads in the circuit.

P4.10 Current Electricity — Circuits

Current electricity is described as movement of charges. It is a particularly useful form of energy because it can be easily transferred from place to place and readily transformed by various devices into other forms of energy (e.g., light, heat, sound, and motion). Electrical current (amperage) in a circuit is determined by the potential difference (voltage) of the power source and the resistance of the loads in the circuit.



P4.10D Discriminate between voltage, resistance, and current as they apply to an electric circuit.



P4.10D Discriminate between voltage, resistance, and current as they apply to an electric circuit.



P4.10D Discriminate between voltage, resistance, and current as they apply to an electric circuit.



P4.10D Discriminate between voltage, resistance, and current as they apply to an electric circuit.



P4.10D Discriminate between voltage, resistance, and current as they apply to an electric circuit.



P4.10D Discriminate between voltage, resistance, and current as they apply to an electric circuit.

P4.10x Current Electricity — Ohm's Law, Work, and Power

In circuits, the relationship between electric current, I , electric potential difference, V , and resistance, R , is quantified by $V = IR$ (Ohm's Law). Work is the amount of energy transferred during an interaction. In electrical systems, work is done when charges are moved through the circuit. Electric power is the amount of work done by an electric current in a unit of time, which can be calculated using $P = IV$.

P4.10x Current Electricity — Ohm's Law, Work, and Power

In circuits, the relationship between electric current, I , electric potential difference, V , and resistance, R , is quantified by $V = IR$ (Ohm's Law). Work is the amount of energy transferred during an interaction. In electrical systems, work is done when charges are moved through the circuit. Electric power is the amount of work done by an electric current in a unit of time, which can be calculated using $P = IV$.

P4.10x Current Electricity — Ohm's Law, Work, and Power

In circuits, the relationship between electric current, I , electric potential difference, V , and resistance, R , is quantified by $V = IR$ (Ohm's Law). Work is the amount of energy transferred during an interaction. In electrical systems, work is done when charges are moved through the circuit. Electric power is the amount of work done by an electric current in a unit of time, which can be calculated using $P = IV$.

P4.10x Current Electricity — Ohm's Law, Work, and Power

In circuits, the relationship between electric current, I , electric potential difference, V , and resistance, R , is quantified by $V = IR$ (Ohm's Law). Work is the amount of energy transferred during an interaction. In electrical systems, work is done when charges are moved through the circuit. Electric power is the amount of work done by an electric current in a unit of time, which can be calculated using $P = IV$.

P4.10x Current Electricity — Ohm's Law, Work, and Power

In circuits, the relationship between electric current, I , electric potential difference, V , and resistance, R , is quantified by $V = IR$ (Ohm's Law). Work is the amount of energy transferred during an interaction. In electrical systems, work is done when charges are moved through the circuit. Electric power is the amount of work done by an electric current in a unit of time, which can be calculated using $P = IV$.

P4.10x Current Electricity — Ohm's Law, Work, and Power

In circuits, the relationship between electric current, I , electric potential difference, V , and resistance, R , is quantified by $V = IR$ (Ohm's Law). Work is the amount of energy transferred during an interaction. In electrical systems, work is done when charges are moved through the circuit. Electric power is the amount of work done by an electric current in a unit of time, which can be calculated using $P = IV$.



P4.10e Explain energy transfer in a circuit, using an electrical charge model.



P4.10e Explain energy transfer in a circuit, using an electrical charge model.



P4.10e Explain energy transfer in a circuit, using an electrical charge model.



P4.10e Explain energy transfer in a circuit, using an electrical charge model.



P4.10e Explain energy transfer in a circuit, using an electrical charge model.



P4.10e Explain energy transfer in a circuit, using an electrical charge model.

P4.10x Current Electricity — Ohm's Law, Work, and Power

In circuits, the relationship between electric current, I , electric potential difference, V , and resistance, R , is quantified by $V = IR$ (Ohm's Law). Work is the amount of energy transferred during an interaction. In electrical systems, work is done when charges are moved through the circuit. Electric power is the amount of work done by an electric current in a unit of time, which can be calculated using $P = IV$.

P4.10x Current Electricity — Ohm's Law, Work, and Power

In circuits, the relationship between electric current, I , electric potential difference, V , and resistance, R , is quantified by $V = IR$ (Ohm's Law). Work is the amount of energy transferred during an interaction. In electrical systems, work is done when charges are moved through the circuit. Electric power is the amount of work done by an electric current in a unit of time, which can be calculated using $P = IV$.

P4.10x Current Electricity — Ohm's Law, Work, and Power

In circuits, the relationship between electric current, I , electric potential difference, V , and resistance, R , is quantified by $V = IR$ (Ohm's Law). Work is the amount of energy transferred during an interaction. In electrical systems, work is done when charges are moved through the circuit. Electric power is the amount of work done by an electric current in a unit of time, which can be calculated using $P = IV$.

P4.10x Current Electricity — Ohm's Law, Work, and Power

In circuits, the relationship between electric current, I , electric potential difference, V , and resistance, R , is quantified by $V = IR$ (Ohm's Law). Work is the amount of energy transferred during an interaction. In electrical systems, work is done when charges are moved through the circuit. Electric power is the amount of work done by an electric current in a unit of time, which can be calculated using $P = IV$.

P4.10x Current Electricity — Ohm's Law, Work, and Power

In circuits, the relationship between electric current, I , electric potential difference, V , and resistance, R , is quantified by $V = IR$ (Ohm's Law). Work is the amount of energy transferred during an interaction. In electrical systems, work is done when charges are moved through the circuit. Electric power is the amount of work done by an electric current in a unit of time, which can be calculated using $P = IV$.

P4.10x Current Electricity — Ohm's Law, Work, and Power

In circuits, the relationship between electric current, I , electric potential difference, V , and resistance, R , is quantified by $V = IR$ (Ohm's Law). Work is the amount of energy transferred during an interaction. In electrical systems, work is done when charges are moved through the circuit. Electric power is the amount of work done by an electric current in a unit of time, which can be calculated using $P = IV$.



P4.10f Calculate the amount of work done when a charge moves through a potential difference, V .



P4.10f Calculate the amount of work done when a charge moves through a potential difference, V .



P4.10f Calculate the amount of work done when a charge moves through a potential difference, V .



P4.10f Calculate the amount of work done when a charge moves through a potential difference, V .



P4.10f Calculate the amount of work done when a charge moves through a potential difference, V .



P4.10f Calculate the amount of work done when a charge moves through a potential difference, V .

P4.10x Current Electricity — Ohm's Law, Work, and Power

In circuits, the relationship between electric current, I , electric potential difference, V , and resistance, R , is quantified by $V = IR$ (Ohm's Law). Work is the amount of energy transferred during an interaction. In electrical systems, work is done when charges are moved through the circuit. Electric power is the amount of work done by an electric current in a unit of time, which can be calculated using $P = IV$.

P4.10x Current Electricity — Ohm's Law, Work, and Power

In circuits, the relationship between electric current, I , electric potential difference, V , and resistance, R , is quantified by $V = IR$ (Ohm's Law). Work is the amount of energy transferred during an interaction. In electrical systems, work is done when charges are moved through the circuit. Electric power is the amount of work done by an electric current in a unit of time, which can be calculated using $P = IV$.

P4.10x Current Electricity — Ohm's Law, Work, and Power

In circuits, the relationship between electric current, I , electric potential difference, V , and resistance, R , is quantified by $V = IR$ (Ohm's Law). Work is the amount of energy transferred during an interaction. In electrical systems, work is done when charges are moved through the circuit. Electric power is the amount of work done by an electric current in a unit of time, which can be calculated using $P = IV$.

P4.10x Current Electricity — Ohm's Law, Work, and Power

In circuits, the relationship between electric current, I , electric potential difference, V , and resistance, R , is quantified by $V = IR$ (Ohm's Law). Work is the amount of energy transferred during an interaction. In electrical systems, work is done when charges are moved through the circuit. Electric power is the amount of work done by an electric current in a unit of time, which can be calculated using $P = IV$.

P4.10x Current Electricity — Ohm's Law, Work, and Power

In circuits, the relationship between electric current, I , electric potential difference, V , and resistance, R , is quantified by $V = IR$ (Ohm's Law). Work is the amount of energy transferred during an interaction. In electrical systems, work is done when charges are moved through the circuit. Electric power is the amount of work done by an electric current in a unit of time, which can be calculated using $P = IV$.

P4.10x Current Electricity — Ohm's Law, Work, and Power

In circuits, the relationship between electric current, I , electric potential difference, V , and resistance, R , is quantified by $V = IR$ (Ohm's Law). Work is the amount of energy transferred during an interaction. In electrical systems, work is done when charges are moved through the circuit. Electric power is the amount of work done by an electric current in a unit of time, which can be calculated using $P = IV$.



P4.10g Compare the currents, voltages, and power in parallel and series circuits.



P4.10g Compare the currents, voltages, and power in parallel and series circuits.



P4.10g Compare the currents, voltages, and power in parallel and series circuits.



P4.10g Compare the currents, voltages, and power in parallel and series circuits.



P4.10g Compare the currents, voltages, and power in parallel and series circuits.



P4.10g Compare the currents, voltages, and power in parallel and series circuits.

P4.10x Current Electricity — Ohm's Law, Work, and Power

In circuits, the relationship between electric current, I , electric potential difference, V , and resistance, R , is quantified by $V = IR$ (Ohm's Law). Work is the amount of energy transferred during an interaction. In electrical systems, work is done when charges are moved through the circuit. Electric power is the amount of work done by an electric current in a unit of time, which can be calculated using $P = IV$.

P4.10x Current Electricity — Ohm's Law, Work, and Power

In circuits, the relationship between electric current, I , electric potential difference, V , and resistance, R , is quantified by $V = IR$ (Ohm's Law). Work is the amount of energy transferred during an interaction. In electrical systems, work is done when charges are moved through the circuit. Electric power is the amount of work done by an electric current in a unit of time, which can be calculated using $P = IV$.

P4.10x Current Electricity — Ohm's Law, Work, and Power

In circuits, the relationship between electric current, I , electric potential difference, V , and resistance, R , is quantified by $V = IR$ (Ohm's Law). Work is the amount of energy transferred during an interaction. In electrical systems, work is done when charges are moved through the circuit. Electric power is the amount of work done by an electric current in a unit of time, which can be calculated using $P = IV$.

P4.10x Current Electricity — Ohm's Law, Work, and Power

In circuits, the relationship between electric current, I , electric potential difference, V , and resistance, R , is quantified by $V = IR$ (Ohm's Law). Work is the amount of energy transferred during an interaction. In electrical systems, work is done when charges are moved through the circuit. Electric power is the amount of work done by an electric current in a unit of time, which can be calculated using $P = IV$.

P4.10x Current Electricity — Ohm's Law, Work, and Power

In circuits, the relationship between electric current, I , electric potential difference, V , and resistance, R , is quantified by $V = IR$ (Ohm's Law). Work is the amount of energy transferred during an interaction. In electrical systems, work is done when charges are moved through the circuit. Electric power is the amount of work done by an electric current in a unit of time, which can be calculated using $P = IV$.

P4.10x Current Electricity — Ohm's Law, Work, and Power

In circuits, the relationship between electric current, I , electric potential difference, V , and resistance, R , is quantified by $V = IR$ (Ohm's Law). Work is the amount of energy transferred during an interaction. In electrical systems, work is done when charges are moved through the circuit. Electric power is the amount of work done by an electric current in a unit of time, which can be calculated using $P = IV$.



P4.10h Explain how circuit breakers and fuses protect household appliances.



P4.10h Explain how circuit breakers and fuses protect household appliances.



P4.10h Explain how circuit breakers and fuses protect household appliances.



P4.10h Explain how circuit breakers and fuses protect household appliances.



P4.10h Explain how circuit breakers and fuses protect household appliances.



P4.10h Explain how circuit breakers and fuses protect household appliances.

P4.10x Current Electricity — Ohm's Law, Work, and Power

In circuits, the relationship between electric current, I , electric potential difference, V , and resistance, R , is quantified by $V = IR$ (Ohm's Law). Work is the amount of energy transferred during an interaction. In electrical systems, work is done when charges are moved through the circuit. Electric power is the amount of work done by an electric current in a unit of time, which can be calculated using $P = IV$.

P4.10x Current Electricity — Ohm's Law, Work, and Power

In circuits, the relationship between electric current, I , electric potential difference, V , and resistance, R , is quantified by $V = IR$ (Ohm's Law). Work is the amount of energy transferred during an interaction. In electrical systems, work is done when charges are moved through the circuit. Electric power is the amount of work done by an electric current in a unit of time, which can be calculated using $P = IV$.

P4.10x Current Electricity — Ohm's Law, Work, and Power

In circuits, the relationship between electric current, I , electric potential difference, V , and resistance, R , is quantified by $V = IR$ (Ohm's Law). Work is the amount of energy transferred during an interaction. In electrical systems, work is done when charges are moved through the circuit. Electric power is the amount of work done by an electric current in a unit of time, which can be calculated using $P = IV$.

P4.10x Current Electricity — Ohm's Law, Work, and Power

In circuits, the relationship between electric current, I , electric potential difference, V , and resistance, R , is quantified by $V = IR$ (Ohm's Law). Work is the amount of energy transferred during an interaction. In electrical systems, work is done when charges are moved through the circuit. Electric power is the amount of work done by an electric current in a unit of time, which can be calculated using $P = IV$.

P4.10x Current Electricity — Ohm's Law, Work, and Power

In circuits, the relationship between electric current, I , electric potential difference, V , and resistance, R , is quantified by $V = IR$ (Ohm's Law). Work is the amount of energy transferred during an interaction. In electrical systems, work is done when charges are moved through the circuit. Electric power is the amount of work done by an electric current in a unit of time, which can be calculated using $P = IV$.

P4.10x Current Electricity — Ohm's Law, Work, and Power

In circuits, the relationship between electric current, I , electric potential difference, V , and resistance, R , is quantified by $V = IR$ (Ohm's Law). Work is the amount of energy transferred during an interaction. In electrical systems, work is done when charges are moved through the circuit. Electric power is the amount of work done by an electric current in a unit of time, which can be calculated using $P = IV$.



P4.10i Compare the energy used in one day by common household appliances (e.g., refrigerator, lamps, hair dryer, toaster, televisions, music players).



P4.10i Compare the energy used in one day by common household appliances (e.g., refrigerator, lamps, hair dryer, toaster, televisions, music players).



P4.10i Compare the energy used in one day by common household appliances (e.g., refrigerator, lamps, hair dryer, toaster, televisions, music players).



P4.10i Compare the energy used in one day by common household appliances (e.g., refrigerator, lamps, hair dryer, toaster, televisions, music players).



P4.10i Compare the energy used in one day by common household appliances (e.g., refrigerator, lamps, hair dryer, toaster, televisions, music players).



P4.10i Compare the energy used in one day by common household appliances (e.g., refrigerator, lamps, hair dryer, toaster, televisions, music players).

P4.10x Current Electricity — Ohm's Law, Work, and Power

In circuits, the relationship between electric current, I , electric potential difference, V , and resistance, R , is quantified by $V = IR$ (Ohm's Law). Work is the amount of energy transferred during an interaction. In electrical systems, work is done when charges are moved through the circuit. Electric power is the amount of work done by an electric current in a unit of time, which can be calculated using $P = IV$.

P4.10x Current Electricity — Ohm's Law, Work, and Power

In circuits, the relationship between electric current, I , electric potential difference, V , and resistance, R , is quantified by $V = IR$ (Ohm's Law). Work is the amount of energy transferred during an interaction. In electrical systems, work is done when charges are moved through the circuit. Electric power is the amount of work done by an electric current in a unit of time, which can be calculated using $P = IV$.

P4.10x Current Electricity — Ohm's Law, Work, and Power

In circuits, the relationship between electric current, I , electric potential difference, V , and resistance, R , is quantified by $V = IR$ (Ohm's Law). Work is the amount of energy transferred during an interaction. In electrical systems, work is done when charges are moved through the circuit. Electric power is the amount of work done by an electric current in a unit of time, which can be calculated using $P = IV$.

P4.10x Current Electricity — Ohm's Law, Work, and Power

In circuits, the relationship between electric current, I , electric potential difference, V , and resistance, R , is quantified by $V = IR$ (Ohm's Law). Work is the amount of energy transferred during an interaction. In electrical systems, work is done when charges are moved through the circuit. Electric power is the amount of work done by an electric current in a unit of time, which can be calculated using $P = IV$.

P4.10x Current Electricity — Ohm's Law, Work, and Power

In circuits, the relationship between electric current, I , electric potential difference, V , and resistance, R , is quantified by $V = IR$ (Ohm's Law). Work is the amount of energy transferred during an interaction. In electrical systems, work is done when charges are moved through the circuit. Electric power is the amount of work done by an electric current in a unit of time, which can be calculated using $P = IV$.

P4.10x Current Electricity — Ohm's Law, Work, and Power

In circuits, the relationship between electric current, I , electric potential difference, V , and resistance, R , is quantified by $V = IR$ (Ohm's Law). Work is the amount of energy transferred during an interaction. In electrical systems, work is done when charges are moved through the circuit. Electric power is the amount of work done by an electric current in a unit of time, which can be calculated using $P = IV$.



P4.10j Explain the difference between electric power and electric energy as used in bills from an electric company.



P4.10j Explain the difference between electric power and electric energy as used in bills from an electric company.



P4.10j Explain the difference between electric power and electric energy as used in bills from an electric company.



P4.10j Explain the difference between electric power and electric energy as used in bills from an electric company.



P4.10j Explain the difference between electric power and electric energy as used in bills from an electric company.



P4.10j Explain the difference between electric power and electric energy as used in bills from an electric company.

P4.1 | x Heat, Temperature, and Efficiency

Heat is often produced as a by-product during energy transformations. This energy is transferred into the surroundings and is not usually recoverable as a useful form of energy. The efficiency of systems is defined as the ratio of the useful energy output to the total energy input. The efficiency of natural and human-made systems varies due to the amount of heat that is not recovered as useful work.

P4.1 | x Heat, Temperature, and Efficiency

Heat is often produced as a by-product during energy transformations. This energy is transferred into the surroundings and is not usually recoverable as a useful form of energy. The efficiency of systems is defined as the ratio of the useful energy output to the total energy input. The efficiency of natural and human-made systems varies due to the amount of heat that is not recovered as useful work.

P4.1 | x Heat, Temperature, and Efficiency

Heat is often produced as a by-product during energy transformations. This energy is transferred into the surroundings and is not usually recoverable as a useful form of energy. The efficiency of systems is defined as the ratio of the useful energy output to the total energy input. The efficiency of natural and human-made systems varies due to the amount of heat that is not recovered as useful work.

P4.1 | x Heat, Temperature, and Efficiency

Heat is often produced as a by-product during energy transformations. This energy is transferred into the surroundings and is not usually recoverable as a useful form of energy. The efficiency of systems is defined as the ratio of the useful energy output to the total energy input. The efficiency of natural and human-made systems varies due to the amount of heat that is not recovered as useful work.

P4.1 | x Heat, Temperature, and Efficiency

Heat is often produced as a by-product during energy transformations. This energy is transferred into the surroundings and is not usually recoverable as a useful form of energy. The efficiency of systems is defined as the ratio of the useful energy output to the total energy input. The efficiency of natural and human-made systems varies due to the amount of heat that is not recovered as useful work.

P4.1 | x Heat, Temperature, and Efficiency

Heat is often produced as a by-product during energy transformations. This energy is transferred into the surroundings and is not usually recoverable as a useful form of energy. The efficiency of systems is defined as the ratio of the useful energy output to the total energy input. The efficiency of natural and human-made systems varies due to the amount of heat that is not recovered as useful work.



P4.11a Calculate the energy lost to surroundings when water in a home water heater is heated from room temperature to the temperature necessary to use in a dishwasher, given the efficiency of the home hot water heater.



P4.11a Calculate the energy lost to surroundings when water in a home water heater is heated from room temperature to the temperature necessary to use in a dishwasher, given the efficiency of the home hot water heater.



P4.11a Calculate the energy lost to surroundings when water in a home water heater is heated from room temperature to the temperature necessary to use in a dishwasher, given the efficiency of the home hot water heater.



P4.11a Calculate the energy lost to surroundings when water in a home water heater is heated from room temperature to the temperature necessary to use in a dishwasher, given the efficiency of the home hot water heater.



P4.11a Calculate the energy lost to surroundings when water in a home water heater is heated from room temperature to the temperature necessary to use in a dishwasher, given the efficiency of the home hot water heater.



P4.11a Calculate the energy lost to surroundings when water in a home water heater is heated from room temperature to the temperature necessary to use in a dishwasher, given the efficiency of the home hot water heater.

P4.1 | x Heat, Temperature, and Efficiency

Heat is often produced as a by-product during energy transformations. This energy is transferred into the surroundings and is not usually recoverable as a useful form of energy. The efficiency of systems is defined as the ratio of the useful energy output to the total energy input. The efficiency of natural and human-made systems varies due to the amount of heat that is not recovered as useful work.

P4.1 | x Heat, Temperature, and Efficiency

Heat is often produced as a by-product during energy transformations. This energy is transferred into the surroundings and is not usually recoverable as a useful form of energy. The efficiency of systems is defined as the ratio of the useful energy output to the total energy input. The efficiency of natural and human-made systems varies due to the amount of heat that is not recovered as useful work.

P4.1 | x Heat, Temperature, and Efficiency

Heat is often produced as a by-product during energy transformations. This energy is transferred into the surroundings and is not usually recoverable as a useful form of energy. The efficiency of systems is defined as the ratio of the useful energy output to the total energy input. The efficiency of natural and human-made systems varies due to the amount of heat that is not recovered as useful work.

P4.1 | x Heat, Temperature, and Efficiency

Heat is often produced as a by-product during energy transformations. This energy is transferred into the surroundings and is not usually recoverable as a useful form of energy. The efficiency of systems is defined as the ratio of the useful energy output to the total energy input. The efficiency of natural and human-made systems varies due to the amount of heat that is not recovered as useful work.

P4.1 | x Heat, Temperature, and Efficiency

Heat is often produced as a by-product during energy transformations. This energy is transferred into the surroundings and is not usually recoverable as a useful form of energy. The efficiency of systems is defined as the ratio of the useful energy output to the total energy input. The efficiency of natural and human-made systems varies due to the amount of heat that is not recovered as useful work.

P4.1 | x Heat, Temperature, and Efficiency

Heat is often produced as a by-product during energy transformations. This energy is transferred into the surroundings and is not usually recoverable as a useful form of energy. The efficiency of systems is defined as the ratio of the useful energy output to the total energy input. The efficiency of natural and human-made systems varies due to the amount of heat that is not recovered as useful work.



P4.11b Calculate the final temperature of two liquids (same or different materials) at the same or different temperatures and masses that are combined.



P4.11b Calculate the final temperature of two liquids (same or different materials) at the same or different temperatures and masses that are combined.



P4.11b Calculate the final temperature of two liquids (same or different materials) at the same or different temperatures and masses that are combined.



P4.11b Calculate the final temperature of two liquids (same or different materials) at the same or different temperatures and masses that are combined.



P4.11b Calculate the final temperature of two liquids (same or different materials) at the same or different temperatures and masses that are combined.



P4.11b Calculate the final temperature of two liquids (same or different materials) at the same or different temperatures and masses that are combined.

P4.12 Nuclear Reactions

Changes in atomic nuclei can occur through three processes: fission, fusion, and radioactive decay. Fission and fusion can convert small amounts of matter into large amounts of energy. Fission is the splitting of a large nucleus into smaller nuclei at extremely high temperature and pressure. Fusion is the combination of smaller nuclei into a large nucleus and is responsible for the energy of the Sun and other stars. Radioactive decay occurs naturally in the Earth's crust (rocks, minerals) and can be used in technological applications (e.g., medical diagnosis and treatment).

P4.12 Nuclear Reactions

Changes in atomic nuclei can occur through three processes: fission, fusion, and radioactive decay. Fission and fusion can convert small amounts of matter into large amounts of energy. Fission is the splitting of a large nucleus into smaller nuclei at extremely high temperature and pressure. Fusion is the combination of smaller nuclei into a large nucleus and is responsible for the energy of the Sun and other stars. Radioactive decay occurs naturally in the Earth's crust (rocks, minerals) and can be used in technological applications (e.g., medical diagnosis and treatment).

P4.12 Nuclear Reactions

Changes in atomic nuclei can occur through three processes: fission, fusion, and radioactive decay. Fission and fusion can convert small amounts of matter into large amounts of energy. Fission is the splitting of a large nucleus into smaller nuclei at extremely high temperature and pressure. Fusion is the combination of smaller nuclei into a large nucleus and is responsible for the energy of the Sun and other stars. Radioactive decay occurs naturally in the Earth's crust (rocks, minerals) and can be used in technological applications (e.g., medical diagnosis and treatment).

P4.12 Nuclear Reactions

Changes in atomic nuclei can occur through three processes: fission, fusion, and radioactive decay. Fission and fusion can convert small amounts of matter into large amounts of energy. Fission is the splitting of a large nucleus into smaller nuclei at extremely high temperature and pressure. Fusion is the combination of smaller nuclei into a large nucleus and is responsible for the energy of the Sun and other stars. Radioactive decay occurs naturally in the Earth's crust (rocks, minerals) and can be used in technological applications (e.g., medical diagnosis and treatment).

P4.12 Nuclear Reactions

Changes in atomic nuclei can occur through three processes: fission, fusion, and radioactive decay. Fission and fusion can convert small amounts of matter into large amounts of energy. Fission is the splitting of a large nucleus into smaller nuclei at extremely high temperature and pressure. Fusion is the combination of smaller nuclei into a large nucleus and is responsible for the energy of the Sun and other stars. Radioactive decay occurs naturally in the Earth's crust (rocks, minerals) and can be used in technological applications (e.g., medical diagnosis and treatment).

P4.12 Nuclear Reactions

Changes in atomic nuclei can occur through three processes: fission, fusion, and radioactive decay. Fission and fusion can convert small amounts of matter into large amounts of energy. Fission is the splitting of a large nucleus into smaller nuclei at extremely high temperature and pressure. Fusion is the combination of smaller nuclei into a large nucleus and is responsible for the energy of the Sun and other stars. Radioactive decay occurs naturally in the Earth's crust (rocks, minerals) and can be used in technological applications (e.g., medical diagnosis and treatment).

E

P4.12A Describe peaceful technological applications of nuclear fission and radioactive decay.

E

P4.12A Describe peaceful technological applications of nuclear fission and radioactive decay.

E

P4.12A Describe peaceful technological applications of nuclear fission and radioactive decay.

E

P4.12A Describe peaceful technological applications of nuclear fission and radioactive decay.

E

P4.12A Describe peaceful technological applications of nuclear fission and radioactive decay.

E

P4.12A Describe peaceful technological applications of nuclear fission and radioactive decay.

P4.12 Nuclear Reactions

Changes in atomic nuclei can occur through three processes: fission, fusion, and radioactive decay. Fission and fusion can convert small amounts of matter into large amounts of energy. Fission is the splitting of a large nucleus into smaller nuclei at extremely high temperature and pressure. Fusion is the combination of smaller nuclei into a large nucleus and is responsible for the energy of the Sun and other stars. Radioactive decay occurs naturally in the Earth's crust (rocks, minerals) and can be used in technological applications (e.g., medical diagnosis and treatment).

P4.12 Nuclear Reactions

Changes in atomic nuclei can occur through three processes: fission, fusion, and radioactive decay. Fission and fusion can convert small amounts of matter into large amounts of energy. Fission is the splitting of a large nucleus into smaller nuclei at extremely high temperature and pressure. Fusion is the combination of smaller nuclei into a large nucleus and is responsible for the energy of the Sun and other stars. Radioactive decay occurs naturally in the Earth's crust (rocks, minerals) and can be used in technological applications (e.g., medical diagnosis and treatment).

P4.12 Nuclear Reactions

Changes in atomic nuclei can occur through three processes: fission, fusion, and radioactive decay. Fission and fusion can convert small amounts of matter into large amounts of energy. Fission is the splitting of a large nucleus into smaller nuclei at extremely high temperature and pressure. Fusion is the combination of smaller nuclei into a large nucleus and is responsible for the energy of the Sun and other stars. Radioactive decay occurs naturally in the Earth's crust (rocks, minerals) and can be used in technological applications (e.g., medical diagnosis and treatment).

P4.12 Nuclear Reactions

Changes in atomic nuclei can occur through three processes: fission, fusion, and radioactive decay. Fission and fusion can convert small amounts of matter into large amounts of energy. Fission is the splitting of a large nucleus into smaller nuclei at extremely high temperature and pressure. Fusion is the combination of smaller nuclei into a large nucleus and is responsible for the energy of the Sun and other stars. Radioactive decay occurs naturally in the Earth's crust (rocks, minerals) and can be used in technological applications (e.g., medical diagnosis and treatment).

P4.12 Nuclear Reactions

Changes in atomic nuclei can occur through three processes: fission, fusion, and radioactive decay. Fission and fusion can convert small amounts of matter into large amounts of energy. Fission is the splitting of a large nucleus into smaller nuclei at extremely high temperature and pressure. Fusion is the combination of smaller nuclei into a large nucleus and is responsible for the energy of the Sun and other stars. Radioactive decay occurs naturally in the Earth's crust (rocks, minerals) and can be used in technological applications (e.g., medical diagnosis and treatment).

P4.12 Nuclear Reactions

Changes in atomic nuclei can occur through three processes: fission, fusion, and radioactive decay. Fission and fusion can convert small amounts of matter into large amounts of energy. Fission is the splitting of a large nucleus into smaller nuclei at extremely high temperature and pressure. Fusion is the combination of smaller nuclei into a large nucleus and is responsible for the energy of the Sun and other stars. Radioactive decay occurs naturally in the Earth's crust (rocks, minerals) and can be used in technological applications (e.g., medical diagnosis and treatment).



P4.12B Describe possible problems caused by exposure to prolonged radioactive decay.



P4.12B Describe possible problems caused by exposure to prolonged radioactive decay.



P4.12B Describe possible problems caused by exposure to prolonged radioactive decay.



P4.12B Describe possible problems caused by exposure to prolonged radioactive decay.



P4.12B Describe possible problems caused by exposure to prolonged radioactive decay.



P4.12B Describe possible problems caused by exposure to prolonged radioactive decay.

P4.12 Nuclear Reactions

Changes in atomic nuclei can occur through three processes: fission, fusion, and radioactive decay. Fission and fusion can convert small amounts of matter into large amounts of energy. Fission is the splitting of a large nucleus into smaller nuclei at extremely high temperature and pressure. Fusion is the combination of smaller nuclei into a large nucleus and is responsible for the energy of the Sun and other stars. Radioactive decay occurs naturally in the Earth's crust (rocks, minerals) and can be used in technological applications (e.g., medical diagnosis and treatment).

P4.12 Nuclear Reactions

Changes in atomic nuclei can occur through three processes: fission, fusion, and radioactive decay. Fission and fusion can convert small amounts of matter into large amounts of energy. Fission is the splitting of a large nucleus into smaller nuclei at extremely high temperature and pressure. Fusion is the combination of smaller nuclei into a large nucleus and is responsible for the energy of the Sun and other stars. Radioactive decay occurs naturally in the Earth's crust (rocks, minerals) and can be used in technological applications (e.g., medical diagnosis and treatment).

P4.12 Nuclear Reactions

Changes in atomic nuclei can occur through three processes: fission, fusion, and radioactive decay. Fission and fusion can convert small amounts of matter into large amounts of energy. Fission is the splitting of a large nucleus into smaller nuclei at extremely high temperature and pressure. Fusion is the combination of smaller nuclei into a large nucleus and is responsible for the energy of the Sun and other stars. Radioactive decay occurs naturally in the Earth's crust (rocks, minerals) and can be used in technological applications (e.g., medical diagnosis and treatment).

P4.12 Nuclear Reactions

Changes in atomic nuclei can occur through three processes: fission, fusion, and radioactive decay. Fission and fusion can convert small amounts of matter into large amounts of energy. Fission is the splitting of a large nucleus into smaller nuclei at extremely high temperature and pressure. Fusion is the combination of smaller nuclei into a large nucleus and is responsible for the energy of the Sun and other stars. Radioactive decay occurs naturally in the Earth's crust (rocks, minerals) and can be used in technological applications (e.g., medical diagnosis and treatment).

P4.12 Nuclear Reactions

Changes in atomic nuclei can occur through three processes: fission, fusion, and radioactive decay. Fission and fusion can convert small amounts of matter into large amounts of energy. Fission is the splitting of a large nucleus into smaller nuclei at extremely high temperature and pressure. Fusion is the combination of smaller nuclei into a large nucleus and is responsible for the energy of the Sun and other stars. Radioactive decay occurs naturally in the Earth's crust (rocks, minerals) and can be used in technological applications (e.g., medical diagnosis and treatment).

P4.12 Nuclear Reactions

Changes in atomic nuclei can occur through three processes: fission, fusion, and radioactive decay. Fission and fusion can convert small amounts of matter into large amounts of energy. Fission is the splitting of a large nucleus into smaller nuclei at extremely high temperature and pressure. Fusion is the combination of smaller nuclei into a large nucleus and is responsible for the energy of the Sun and other stars. Radioactive decay occurs naturally in the Earth's crust (rocks, minerals) and can be used in technological applications (e.g., medical diagnosis and treatment).

E

P4.12C Explain how stars, including our Sun, produce huge amounts of energy (e.g., visible, infrared, ultraviolet light).

E

P4.12C Explain how stars, including our Sun, produce huge amounts of energy (e.g., visible, infrared, ultraviolet light).

E

P4.12C Explain how stars, including our Sun, produce huge amounts of energy (e.g., visible, infrared, ultraviolet light).

E

P4.12C Explain how stars, including our Sun, produce huge amounts of energy (e.g., visible, infrared, ultraviolet light).

E

P4.12C Explain how stars, including our Sun, produce huge amounts of energy (e.g., visible, infrared, ultraviolet light).

E

P4.12C Explain how stars, including our Sun, produce huge amounts of energy (e.g., visible, infrared, ultraviolet light).

P4.12x Mass and Energy

In nuclear reactions, a small amount of mass is converted to a large amount of energy, $E = mc^2$, where c is the speed of light in a vacuum. The amount of energy before and after nuclear reactions must consider mass changes as part of the energy transformation.

P4.12x Mass and Energy

In nuclear reactions, a small amount of mass is converted to a large amount of energy, $E = mc^2$, where c is the speed of light in a vacuum. The amount of energy before and after nuclear reactions must consider mass changes as part of the energy transformation.

P4.12x Mass and Energy

In nuclear reactions, a small amount of mass is converted to a large amount of energy, $E = mc^2$, where c is the speed of light in a vacuum. The amount of energy before and after nuclear reactions must consider mass changes as part of the energy transformation.

P4.12x Mass and Energy

In nuclear reactions, a small amount of mass is converted to a large amount of energy, $E = mc^2$, where c is the speed of light in a vacuum. The amount of energy before and after nuclear reactions must consider mass changes as part of the energy transformation.

P4.12x Mass and Energy

In nuclear reactions, a small amount of mass is converted to a large amount of energy, $E = mc^2$, where c is the speed of light in a vacuum. The amount of energy before and after nuclear reactions must consider mass changes as part of the energy transformation.

P4.12x Mass and Energy

In nuclear reactions, a small amount of mass is converted to a large amount of energy, $E = mc^2$, where c is the speed of light in a vacuum. The amount of energy before and after nuclear reactions must consider mass changes as part of the energy transformation.



P4.12d Identify the source of energy in fission and fusion nuclear reactions.



P4.12d Identify the source of energy in fission and fusion nuclear reactions.



P4.12d Identify the source of energy in fission and fusion nuclear reactions.



P4.12d Identify the source of energy in fission and fusion nuclear reactions.



P4.12d Identify the source of energy in fission and fusion nuclear reactions.



P4.12d Identify the source of energy in fission and fusion nuclear reactions.